

01-21-00

A

PTO/SB/05 (12/97)

Approved for use through 09/30/00. OMB 0651-0032

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number

UTILITY PATENT APPLICATION TRANSMITTAL <i>(Only for new nonprovisional application under 37 CFR 1.53(b))</i>		CERTIFICATE UNDER 37 C.F.R. 1.10(b)		Docket No. A/B-1006 Div-IV
		Express Mail Label No.: EI743163685US		Date of Deposit: Jan. 20, 2000
APPLICATION ELEMENTS <i>See MPEP chapter 600 concerning utility patent application contents.</i>		I hereby certify that this paper is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" Service on the date indicated above, and is address to: Asst. Commissioner for Patents, Box Patent Application, Washington, DC 20231		
		Signature and Date <i>Imre Balogh Jan. 20/2000</i>		
ADDRESS TO: Assistant Commissioner for Patents Box Patent Application Washington, DC 20231				
1. <input checked="" type="checkbox"/> Fee Transmittal Form <i>(Submit an original, and a duplicate for fee processing)</i> 2. <input checked="" type="checkbox"/> Specification <i>Total Pages</i> 78 <i>(preferred arrangement set forth below)</i> <ul style="list-style-type: none"> - Descriptive Title of the Invention - Cross References to Related Applications - Statement Regarding Fed sponsored R & D - Reference to Microfiche Appendix - Background of the Invention - Brief Summary of the Invention - Brief Description of the Drawings <i>(if filed)</i> - Detailed Description - Claim(s) - Abstract of the Disclosure 3. <input type="checkbox"/> Drawing(s) <i>(35 USC 113)</i> <i>Total Sheets</i> <input type="checkbox"/> 4. <input checked="" type="checkbox"/> Oath or Declaration/ <i>Total Sheets</i> 3 Power of Attorney <ul style="list-style-type: none"> a. <input type="checkbox"/> Newly executed (original or copy) b. <input checked="" type="checkbox"/> Copy from a prior application (37 CFR 1.63(d)) <i>(for continuation/divisional with Box 17 completed - [Note Box 5 below])</i> i. <input type="checkbox"/> DELETION OF INVENTOR(S) Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b). 5. <input checked="" type="checkbox"/> Incorporation By Reference <i>(useable if Box 4b is checked)</i> The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.				
6. <input type="checkbox"/> Microfiche Computer Program <i>(Appendix)</i> 7. <input type="checkbox"/> Nucleotide and/or Amino Acid Sequence Submission <i>(if applicable, all necessary)</i> <ul style="list-style-type: none"> a. <input type="checkbox"/> Computer Readable Copy b. <input type="checkbox"/> Paper copy (identical to computer copy) c. <input type="checkbox"/> Statement verifying identity of above copies 				
ACCOMPANYING APPLICATION PARTS				
8. <input type="checkbox"/> Assignment Papers (cover sheet & document(s)) 9. <input type="checkbox"/> 37 CFR 3.73(b) Statement <input type="checkbox"/> Power of Attorney 10. <input type="checkbox"/> English Translation Document <i>(if applicable)</i> 11. <input checked="" type="checkbox"/> Information Disclosure <input type="checkbox"/> Copies of IDS Statement (IDS)/PTO- Citations 1449 12. <input checked="" type="checkbox"/> Preliminary Amendment 13. <input checked="" type="checkbox"/> Return Receipt Postcard (MPEP 503) <i>(Should be specifically itemized)</i> 14. <input checked="" type="checkbox"/> Small Entity <input checked="" type="checkbox"/> Statement filed in prior application Statement(s) Status still proper and desired 15. <input type="checkbox"/> Certified Copy of Priority Document(s) <i>(if foreign priority is claimed)</i> 16. <input checked="" type="checkbox"/> Other <input type="checkbox"/> Check - \$345.00				
17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information: <input type="checkbox"/> Continuation <input checked="" type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-part (CIP) of prior application no: 09/411,111				
18. CORRESPONDENCE ADDRESS				
The Law Offices of Imre Balogh 276 Smith School Road Perkasi, PA 18944 Telephone No.: (215) 249-9287 Facsimile No.: (215) 249-1286			Respectfully submitted, <i>Imre Balogh</i> IMRE BALOGH Reg. No. 26,835	

 01/20/00
 10654 U.S. PTO

 09/488420
 01/20/00

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: Farrar et al.

Prior Application: 09/411,111

Filing Date: October 4, 1999

Anticipated Classification of this Application:

Class: 514 Subclass: 150

Art Unit: 1614

Examiner: V. Kim

For: Kappa Agonist Anti-Pruritic
Pharmaceutical Formulations and Method of
Treating Pruritus Therewith

Assistant Commissioner for Patents
Washington, D.C. 20231

Dear Sir:


CERTIFICATE UNDER 37 C.F.R. 1.10(b)

Express Mail Label Number: EI743163685US

Date of Deposit: Jan. 20, 2000

I hereby certify that this paper is being deposited with the
United States Postal Service "Express Mail Post Office to
Addressee" Service on the date indicated above and is
addressed to: Asst. Commissioner for Patents,
Washington, DC 20231

Signature



PRELIMINARY AMENDMENT

This is a Preliminary Amendment of a divisional of co-pending U.S. Application Serial No. 09/411,111 filed on October 4, 1999.

Please amend the above-identified application as requested herein.

IN THE SPECIFICATION

Please delete the continuing application data on page 1 and replace it with the following:


-- This application is a divisional of Application Ser. No. 09/411,111 filed on October 4, 1999 which in turn is a divisional of Application Ser. No. 09/184,393 filed on November 2, 1998, now U.S. Patent No. 6,004,694 which in turn is a divisional of Application Ser. No. 08/064,695 filed on April 22, 1998, now U.S. Patent No. 5,869,521, which in turn is a divisional of Application Ser. No. 08/892,599 filed on July 14, 1997, now U.S. Patent No. 5,760,023.--

IN THE CLAIMS

Please cancel claims 1-62.

Please amend the following claims as shown:.

--67. (Amended) A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition [of claim 63.] formula V or a pharmaceutically acceptable salt thereof



wherein

----- represents a single or double bond;

R¹ represents an alkyl group having 1-5 carbon atoms, a cycloalkylalkyl group having 4-7 carbon atoms, a cycloalkenylalkyl group having 5-7 carbon atoms, an aryl group having 6-12 carbon atoms, an aralkyl group having 7-13 carbon atoms, an alkenyl group having 4-7 carbon atoms, an allyl group, a furan-2-ylalkyl group having 1-5 carbon atoms, or a thiophen-2-ylalkyl group having 1-5 carbon atoms;

R² represents a hydrogen atom, a hydroxy group, a nitro group, an alkanoyloxy group having 1-5 carbon atoms, an alkoxy group having 1-5 carbon atoms, an alkyl group having 1-5 carbon atoms, or -NR⁹R¹⁰ wherein R⁹ represents a hydrogen atom or an alkyl group having 1-5 carbon atoms, and R¹⁰ represents a hydrogen atom, an alkyl group having 1-5 carbon atoms, or -C(=O)R¹¹

wherein R¹¹ represents a hydrogen atom, a phenyl group or an alkyl group having 1-5 carbon atoms;

R³ represents a hydrogen atom, a hydroxy group, an alkanoyloxy group having 1-5 carbon atoms, or an alkoxy group having 1-5 carbon atoms;











A represents -XC(=Y)-, -XC(=Y)Z-, -X-, -XSO₂-, or -OC(OR⁴)R⁴- where, X, Y and Z each independently represent NR⁴, S or O wherein R⁴ represents a hydrogen atom, a straight-chain or branched chain alkyl group having 1-5 carbon atoms or an aryl group having 6-12 carbon atoms, and wherein R⁴ may be identical or different;

B represents a valence bond, a straight-chain or branched chain alkylene group having 1-14 carbon atoms which may be substituted with at least one substituent selected from the group consisting of an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, a hydroxy group, fluorine, chlorine, bromine, iodine, an amino group, a nitro group, a cyano group, a trifluoromethyl group and a phenoxy group, and wherein 1 to 3 methylene groups may be replaced with carbonyl groups, an acyclic unsaturated hydrocarbon containing from 1 to 3 double bonds and/or triple bonds and having 2-14 carbon atoms which may be substituted with at least one substituent group selected from the group consisting of an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, a hydroxy group, fluorine, chlorine, bromine, iodine, an amino group, a nitro group, a cyano group, a trifluoromethyl group and a phenoxy group, and wherein from 1 to 3 methylene groups may be replaced with carbonyl groups, or a straight-chain or branched chain saturated or unsaturated hydrocarbon group containing from 1 to 5 thioether, ether and/or amino bonds and having 1-14 carbon atoms wherein hetero atoms are not bonded directly to A, and 1 to 3 methylene groups may be replaced with carbonyl groups;

R⁵ represents a hydrogen atom or an organic group which may be substituted with at least one or more substituent groups selected from the group consisting of an alkyl group having 1-5 carbon atoms, an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms,

a hydroxy group, fluorine, chlorine, bromine, iodine, an amino group, a nitro group, a cyano group, an isothiocyanate group, a trifluoromethyl group and a methylenedioxy group; or

R_5 is



wherein

Q is N, O or S;

T is CH, N, S or O;

l is 0-5;

m and n are > 0

$m + n < 5$;

R^6 represents a hydrogen atom;

R^7 represents a hydrogen atom, a hydroxy group, an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, or R^6 and R^7 together represent -O-, -CH₂- or -S-;
and


R⁸ represents a hydrogen atom, an alkyl group having 1-5 carbon atoms, or an alkanoyl group having 1-5 carbon atoms in a pharmaceutically acceptable carrier.--

--72. (Amended) A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition of claim [64.] 67 wherein

R¹ is an alkyl group having 1-5 carbon atoms, a cycloalkylmethyl group having 4-7 carbon atoms, a cycloalkenylmethyl group having 5-7 carbon atoms, a phenylalkyl group having 7-13 carbon atoms, an alkenyl group having 4-7 carbon atoms, an allyl group, a furan-2-yl-alkyl group having 1-5 carbon atoms and a thiophen-2-yl-alkyl group having 1-5 carbon atoms;

R² is hydrogen, hydroxy, nitro, acetoxy, methoxy, methyl, ethyl, propyl, amino, dimethylamino, acetylamino or benzoylamino groups; or

R⁴ is



Formula V-1

wherein

Q is N, O or S;

T is CH, N, S or O;

m and n are > 0 and

m + n < 5;

B is $-(CH_2)_n-$ wherein $n = 0-6$, $-(CH_2)_n-C(=O)-$ wherein $n = 1-4$, $-CH=CH-(CH_2)_n-$ wherein $n = 0-4$, $-C\equiv C-(CH_2)_n-$ wherein $n=0-4$, $-CH_2-O-$, $-CH_2-S-$, $-CH_2-O-(CH_2)_2-O-(CH_2)_2-$, $-CH_2-O-CH_2-NH-CH_2-O-CH_2-$ and $-CH_2-O-CH_2-S-CH_2-O-CH_2-$;

R^5 is hydrogen or an organic group of Formula V-1 said organic group may be substituted with at least one substituent group selected from the group consisting of an alkyl group having 1-5 carbon atoms, an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, a hydroxy group, fluorine, chlorine, bromine, an amino group, a nitro group, a cyano group, an isothiocyanate group and a trifluoromethyl group.--

--77. (Amended) A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition of claim [65.] 72 wherein

R^1 is methyl, ethyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclopentenylmethyl, cyclohexenylmethyl, benzyl, phenethyl, trans-2-butenyl, 2-methyl-2-butenyl, allyl, furan-2-yl-methyl or thiophen-2-yl-methyl;

R^2 is hydrogen, hydroxy, nitro, acetoxy, methyl or dimethylamino;

R^3 is $-NR^4C(=O)-$, $-NR^4C(=S)-$, $-NR^4C(=O)O-$, $-NR^4C(=O)NR^4-$, $-NR^4C(=S)NR^4-$ or $-NR^4SO_2-$;

R^4 is a straight-chain or branched alkyl group having 1-5 carbon atoms;

B is $-(CH_2)_n-$ wherein $n=0-6$, $-CH=CH(CH_2)_n-$ wherein $n=0-4$, $-C\equiv C-(CH_2)_n-$ wherein $n=0-4$, $-CH_2-O-$ or $-CH_2-S-$; and

R^5 is hydrogen, phenyl, 3,4-dichlorophenyl, 4-chlorophenyl, 3-chlorophenyl, 3,4-difluorophenyl, 4-fluorophenyl, 3-fluorophenyl, 2-fluorophenyl, 4bromophenyl, 3-bromophenyl, 2-bromophenyl, 4-nitrophenyl, 3-nitrophenyl, 2-nitrophenyl, 4-trifluoromethylphenyl, 3-trifluoromethylphenyl, 2-trifluoromethylphenyl, 4-methylphenyl, 3-methylphenyl, 2-methylphenyl, 4-methoxyphenyl, 3-methoxyphenyl, 2-methoxy, 3-furanyl, 2-furanyl, 3-thienyl, 2-thienyl, cyclopentyl or cyclohexyl--

--82. (Amended) A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition of claim [66.] 77 wherein said compound is elected from the group consisting of:

17-cyclopropylmethyl-4,5a-epoxy-3,14b-dihydroxy-6b-(N-methyl-3-phenylpropionamido)morphinan;

17-cyclopropylmethyl-4,5a-epoxy-3,14b-dihydroxy-6b-(N-methyl-trans-3-(3-furyl)acrylamido)morphinan;

17-cyclopropylmethyl-4,5a-epoxy-3,14b-dihydroxy-6b-(N-methyl-trans-3-cyclohexylacrylamido)morphinan;

17-cyclopropylmethyl-4,5a-epoxy-3,14b-dihydroxy-6b-(N-methyl-trans-3-(4-trifluoromethylphenyl)acrylamido)morphinan;

17-cyclopropylmethyl-4,5a-epoxy-3,14b-dihydroxy-6a-(N-methyl-trans-3-(3-thiophenyl)acrylamido)morphinan;

17-cyclopropylmethyl-4,5a-epoxy-3,14b-dihydroxy-6b-(N-methyl-trans-3-phenylacrylamido)morphinan;

17-cyclopropylmethyl-4,5a-epoxy-3,14b-dihydroxy-6b-(N-methyl-trans-2-hexenamido)morphinan; and

17-cyclopropylmethyl-4,5a-epoxy-3,14b-dihydroxy-6b-(N-methyl-phenylpropiolamido)morphinan.--

REMARKS

In parent application Serial No. 08/892,599 (now U.S. Patent No. 5,760,023) claims 1-18 were prosecuted covering Formula I.

In the first divisional application Serial No. 08/064,695 claims 19-38 were prosecuted covering formulas II, IIa and IIb (now U.S. Patent No. 5,869,521).

In the second divisional application Serial No. 09/184,393, claims 39-50 were prosecuted covering formula III. This case has now issued as U.S. Patent No. 6,004,694.

In the third divisional application Serial No. 09/411,111, claims 51-62 were prosecuted covering formula IV. This case is now allowed.

In the present (fourth) divisional application, the remaining claims 63-86 covering formula V is wished, and respectfully requested, by the applicants to be prosecuted.

In the parent application and in the first, second and third divisional applications, applicants have canceled the composition claims since publications/patents disclose the majority of compounds/compositions having kappa opioid agonist activity. However, none of the publications/patents disclose or suggest the use of the compounds/compositions for the prevention or treatment of pruritus.

In the present, fourth divisional application, applicants have also canceled the composition claims and amended the method of treatment claims.

As a result of this preliminary amendment:

Claims 1-66 are canceled;


Claims 67, 72, 77 and 82 are amended; and

Claims 68-71, 73-76, 78-81 and 83-86 remain as originally filed.

Favorable consideration of the application is respectfully requested.

Respectfully submitted,

Date Jan. 20, 2000



Imre Balogh
Reg. No. 26,835

Address:
276 Smith School Road
Perkasie, PA 18944
Tele: (215) 249-9287
Facsimile: (215) 249-1286

**KAPPA AGONIST ANTI-PRURITIC PHARMACEUTICAL FORMULATIONS
AND METHOD OF TREATING PRURITUS THEREWITH**

BACKGROUND OF THE INVENTION

This application is a divisional of Application Ser. No. 09/064,695 filed on April 22, 1998, which in turn is a divisional of Application Ser. No. 08/892,599 filed on July 14, 1997, now U.S. Patent No. 5,760,023.--

1. Field of the Invention

This invention relates to pharmaceutical formulations containing kappa agonist compounds and to their medical use as anti-pruritic agonist at kappa opioid receptors.

2. Reported Developments

A) Antihyperalgesic Kappa Agonists

Opium and its derivatives are potent analgesics that also have other pharmacological effects, and exert their effects by interacting with high-affinity receptors.

It has been shown by investigators that there are at least three major opioid receptor types in the central nervous system (hereinafter CNS) and in the periphery. These receptors, known as mu (μ), delta (δ) and kappa (κ), have distinct pharmacological profiles, anatomical distributions and functions. [See, for example: Wood, P.L., Neuropharmacology, 21, 487-497, 1982; Simon, E. J., Med. Res. Rev., 11, 357-374, 1991; Lutz et al, J. Recept. Res. 12, 267-286; and Mansour et al, Opioid I, ed. Herz, A. (Springer, Berlin) pp. 79-106, 1993.] The δ receptors are abundant in CNS and mediate analgesia, gastrointestinal motility and various hormonal functions. The μ receptors bind morphine-like drugs and mediate the opiate phenomena associated with morphine, including analgesia, opiate dependence, cardiovascular and respiratory functions, and several neuroendocrine effects.

The κ receptors have a wide distribution in CNS and mediate a spectrum of functions including the modulation of drinking, water balance, food intake, gut motility, temperature control and various endocrine functions. They also produce analgesia. [See, for example: Leander et al, J. Pharmacol. Exp. Ther. 234, 463-469, 1985; Morley et al, Peptides 4, 797-800, 1983; Manzanares et al, Neuroendocrinology 52, 200-205, 1990; and Iyengar et al, J. Pharmacol. Exp. Ther, 238, 429-436, 1986.]

Most clinically used opioid analgesics such as morphine and codeine act as μ receptor agonists. These opioids have well-known, undesirable and potentially dangerous dependence forming side effects. Compounds which are κ -receptor agonists act as analgesics through interaction with κ opioid receptors. The advantage of these agonists over the classical μ receptor agonists, such as morphine, lies in their ability to cause analgesia while being devoid of morphine-like behavioral effects and addiction liability.

B) Antihyperalgesic Kappa Agonists as Anti-Pruritic Agents

The prior art has investigated the physiology and treatment of pruritus as illustrated hereunder.

Itch is a well known sensory state associated with the desire to scratch. As with pain, itch can be produced by a variety of chemical, mechanical, thermal or electrical stimuli. In addition to the difference in the sensory quality of itch and pain, they also differ in that (1) itch, unlike pain, can only be evoked from the superficial layers of skin, mucosa, and conjunctiva, and (2) itch and pain usually do not occur simultaneously from the same skin region; in fact, mildly painful stimuli, such as scratching, are effective in eliminating itch. In addition, the application of histamine to skin produces itch but not pain. Itch and pain are further dissociated pharmacologically: itch appears to be insensitive to opiate and non-steroidal anti-inflammatory drug (NSAID) treatment, both of which are effective in treating pain.

Although itch and pain are of a class in that both are modalities of nociception transmitted by small unmyelinated C fibers, evidence that itch is not just a variety of low-threshold pain is overwhelming. Itch leads to the reflex or urge to scratch; pain leads to withdrawal. Removal of the epidermis eliminates itch but causes pain. Analgesics, particularly opioids, relieve pain but often cause itch (see, for example *J. Am. Acad. Derm.* 24: 309-310, 1991). There can be no doubt that itching is of eminent clinical importance; many systemic and skin diseases are accompanied by persistent or recurrent itch attacks. Current knowledge suggests that itch has several features in common with pain but exhibits intriguing differences as well (see, for example, W. Magerl, *IASP Newsletter*, pp. 4-7, Sept/Oct 1996).

McMahon et al (*TINS*, Vol. 15, No. 12, pp. 497-501, 1992) provides a description of stimuli (Table a) and a comparison of the established features of itch and pain (Table b):

Table a
Stimuli that can elicit or augment itch

5	Physical	<p><i>Mechanical.</i> Light touch, pressure, suction.</p> <p><i>Thermal.</i> Warming.</p> <p><i>Electrical.</i> Focal transcutaneous repetitive stimulation, transcutaneous constant current stimulation, intraneural microstimulation.</p>
10	Chemical	<p><i>Non-specific irritants.</i> Acids, alkalis.</p> <p><i>Inflammatory mediators.</i> Histamine, kallikrein, bradykinin, prostaglandins.</p> <p><i>Histamine-releasing substances.</i> Compound 48/80, protamine, C3a.</p>
15		<p><i>Peptidases.</i> Mucinain, papain, trypsin, mast cell chymase.</p> <p><i>Neuropeptides.</i> Substance P, vasoactive intestinal polypeptide, neurotensin, secretin.</p> <p><i>Opioids.</i> Morphine, β-endorphin, enkephalin analogues.</p>

Table b
Comparison of the established features of itch and pain

	ITCH	PAIN
<i>Psychophysiology</i>		
Tissue	Skin. Mucous membranes	Most tissues
Stimulus	See Table a	Many stimuli
Intraneural microstimulation	Occasionally	Yes
Secondary sensations	Allokinesis (itchy skin)	Hyperalgesia
Psychogenic modification	Pronounced	Present
Counterstimuli	Scratching, pain, cooling	Tactile stimuli, cooling
<i>Neurophysiology</i>		
Primary afferent neurones	C- and A δ -fibres	C- and A δ -fibres
Flare size	Large	Small
Spinal pathway	Anterolateral funiculus	Anterolateral funiculus
Protective reflexes	Scratching, sneezing	Flexion, guarding
Autonomic reflexes	Yes	Yes
<i>Pharmacology</i>		
Capsaicin sensitivity	Yes	Chemogenic pain; yes
NSAID sensitivity	Probably not	Yes
Morphine sensitivity	No	Yes

Abbreviation: NSAID, non-steroidal anti-inflammatory drugs.

5

Experimental focal itch stimuli are surrounded by a halo of seemingly unaffected tissue where light tactile stimuli are capable of eliciting itch-like sensations. The term itchy skin or allokinesis has been coined for these secondary sensations that are reminiscent of the features of secondary hyperalgesia evolving around a painful focus. A crucial observation is that itch and pain usually do not coexist in the same skin region and a mild noxious stimulus such as scratching is in fact the singly most effective way to abolish itch. This abolition of itch can be prolonged producing an 'antipruritic state'. Although mild scratch is often not painful, microneurographic recordings from humans have directly determined that such stimuli are among the most effective ways to excite cutaneous unmyelinated nociceptive afferents. (See, for example:

15

Shelly, W.B. and Arthur, R.P. (1957) *Arch. Dermatol.* 76, 296-323;

Simone, D.A. et al. (1987) *Somatosens. Res.* 5, 81-92;

Graham, D.T. , Goodell, H. and Wolff, H.G. (1951) *J. Clin. Invest.* 30, 37-49;

Simone, D.A., Alreja, M. and LaMotte, R.H. (1991) *Somatosens, Mot. Res.* 8, 271-279;

Torebjörk, E (1985) *Philos. Trans. R. Soc. London Ser. B* 308, 227-234; and

Vallbo, A.B., Hagbarth, K.E., Torebjörk, H.E. and Wallin, B.G. (1979) *Physiol. Rev.* 59, 919-957).

Physiologically, there is evidence that substance P released from nociceptor terminals can cause the release of histamine from mast cells. Activation of mast cells, with release of the pruritogen histamine, occurs in immediate type hypersensitivity diseases, such as anaphylactic reactions and urticaria. Urticarial eruptions are distinctly pruritic and can involve any portion of the body, and have a variety of causes beyond hypersensitivity, including physical stimuli such as cold, solar radiation, exercise and mechanical irritation. Other causes of pruritus include: chiggers, the larval form of which secretes substance that creates a red papule that itches intensely; secondary hyperparathyroidism associated with chronic renal failure; cutaneous larva migrans, caused by burrowing larvae of animal hookworms; dermal myiasis, caused by maggots of the horse botfly, which can afflict horseback riders; onchocerciasis ("river blindness") caused by filarial nematodes; pediculosis, caused by lice infestations; enterobiasis (pinworm) infestations, which afflict about 40 million Americans, particularly school children; schistosome dermatitis (swimmer's itch); psoriasis; poison ivy; and asteatotic eczema ("winter itch"). The role of histamine or other endogenous pruritogens in mediating itch associated with these and other pruritic conditions, such as atopic dermatitis, is not yet well established. For atopic dermatitis, in particular, it appears that itch is not inhibited by antihistamines, but by cyclosporin A, a drug which inhibits the production of cytokines which have been proposed as potential pruritogens.

Current therapies for the treatment of itch include a variety of topical and systemic agents, such as steroids, antihistamines, and some psychotherapeutic tricyclic compounds, such as doxepin hydrochloride. Many such agents are listed in *PDR Generics* (see Second Edition, 1996, p. cv for a listing of said agents). The limitations of these agents are well known to medical practitioners, and are summarized in the "Warnings" and "Precautions" sections for the individual agents listed in *PDR Generics*. In particular, the lack of complete efficacy of antihistamines is well known, but antihistamines are frequently used in dermatology to treat pruritus due to urticaria, atopic dermatitis, contact dermatitis, psoriasis, and a variety of other conditions. Although sedation has been a frequent side effect of conventional systemically administered antihistamines, a new generation of antihistamines have been developed that are non-sedating, apparently due to their inability to cross the blood-brain barrier.

Intravenous administration of opiate analgesics, such as morphine and hydromorphone has been associated with pruritus, urticaria, other skin rashes, and wheal and flare over the vein being injected. These itch and itch-related reactions are believed to be due to a histamine-releasing property of these opiates, via mast cell degranulation.

5 These opiates are thought to act upon the mu subtype of opiate receptor, but the possibility of interactions at the other principal opiate receptor subtypes (delta and kappa) cannot be excluded since these and other pruritogenic analgesics are not pure mu agonists. The cellular loci of the receptor type(s) mediating the itching effect is not known, although the mast cell is a possible candidate since opiates cause histamine
10 release from these cells. However, some investigators have suggested that the frequent inability of antihistamines to block morphine-induced itching suggests a non-histaminergic mediation of opiate-induced itching - mechanism which could involve central opiate receptors. Although i.v. morphine only occasionally results in generalized itching (in about 1% of patients), pruritus is more prevalent in opiate analgesia with
15 epidural (8.5%) or intraspinal (45.8%) administration. (See, for example: Bernstein et al., "Antipruritic Effect of an Opiate Antagonist, Naloxone Hydrochloride", *The Journal of Investigative Dermatology*, 78:82-83, 1982; and Ballantyne et al., "Itching after epidural and spinal opiates", *Pain*, 33: 149-160, 1988.)

To date, treatment with opiates has not only proven useless in the treatment of
20 itch, but appears to exacerbate itch in mammals. The consistent findings from human studies indicate that whether by central or peripheral mechanisms, opiates appear to promote rather than prevent itching, and that opiate antagonists have antipruritic activity.

Human clinical studies have generally shown that opiates cause itching and there is evidence that these effects can be reproduced in animal models, where itching
25 sensations per se cannot be reported, but scratching behavior can be observed. (See, for example: Thomas et al., "Microinjection of morphine into the rat medullary dorsal horn produces a dose-dependent increase in facial-scratching", *Brain Research*, 695: 267-270, 1996; Thomas et al., "Effects of central administration of opioids on facial scratching in monkeys", *Brain Res.*, 585: 315-317, 1992; and Thomas et al., "The medullary dorsal
30 horn: A site of action of opioids in producing facial scratching in monkeys", *Anesthesiology*, 79: 548-554, 1993).

We have now discovered that certain kappa agonists, which are substantially devoid of central nervous system effects, possess anti-pruritic activity in addition to antihyperalgesic activity. Accordingly, the present invention also provides safe and
35 effective compositions for the prevention and treatment of pruritus.


Compounds having kappa opioid agonist activity are disclosed in the following references all of which are incorporated herein by reference:

1. U.S. Patent No. 4,145,435
- 5 2. U. S. Patent No. 4,360,531
3. U. S. Patent No. 4,359,476
4. EPA 0 108 602
5. U. S. Patent No. 4,855,316
6. EPA 0 393 696
- 10 7. EPA 0 372 466
8. U.S. Patent No. 4,906,655
9. U.S. Patent No. 4,438,130
10. U.S. Patent No. 4,663,343
11. U. S. Patent No. 5,114,945
- 15 12. U.S. Patent No. 4,943,578
13. EPA 0 330 467
14. EPA 0 366 327
15. EPA 0 398 720
16. EPA 0 330 469
- 20 17. WO 92/20657
18. EPA 0 409 489 A2
19. EPA 0 333 427
20. WO 90/07502
21. EPA 0 356 247
- 25 22. EP 0 752 246 A2
23. WO 96/06077
24. EP 0 483 580
25. EPA 0 254 545
26. EPA 0 325 406
- 30 27. EPA 0 261 842
28. WO 96/06077
29. WO 94/18165
30. WO 96/06078
31. EPA 0 577 847 A1.
- 35 32. U.S. Patent No. 4,929,627 and
33. EPA 0 260 555

SUMMARY OF THE INVENTION

Compositions containing compounds having kappa opioid agonist activity are provided for preventing or treating pruritus. A compound of formulae I, II, IIa, IIb, III, IV and V, or a pharmaceutically acceptable salt thereof, are used in the compositions.

The compounds of formula I have the following structure:

**I**

10 wherein

the wavy line bond (~) between the nitrogen in the 2-position and the cyclohexyl ring carbon indicates the bond can be either cis- or trans with respect to each substituent on the cyclohexyl ring;

15

A is a single chemical bond (-), $-(CH_2)_q$, $CH(CH_3)-$ or $-X(CH_2)_n$
 where q is 1 to 4,
 n is 1-4 and
 x is O or S;

20

Ar is an aromatic, hetero-aromatic, bicyclic-aromatic, tricyclic-aromatic group or diphenyl methyl each of which may be unsubstituted or substituted with a member selected from the group consisting of H, halo, trifluoromethyl, nitro, C_1-C_3 -alkoxy, hydroxy, azido, C_1-C_3 -alkyl, methanesulfonyl, cyano, amino, C_1-C_3 -alkoxycarbonyl, C_1-C_3 -alkanoyloxy, and C_1-C_3 -carboxacylamino of the formula $-NHC(O)R_7$

25

where R_7 is H, C_1-C_2 -alkyl, or an aromatic or hetero-aromatic group;

R_1 and R_2 are independently H, C_1-C_3 -alkyl or allyl;

30


R_1 and R_2 , taken together with the nitrogen to which they are bonded, complete a ring selected from the group consisting of azetidiny, pyrrolidiny, 3-hydroxypyrrolidiny, 3-fluoropyrrolidiny, morpholiny, piperidiny, and 3,4-dehydropiperidiny;

R_3 , R_4 , R_5 , R_6 are independently H, hydroxy, OR_8 or $OC(=O)R_9$;

R_5 and R_6 taken together may form the group $-E-CH_2-CH_2-E-$;

R_5 and R_6 taken together form a ring

5



where

Z is selected from the group consisting of oxygen (-O-), NR_{10} , sulfur (-S-), sulfinyl (-S(O)-), and sulfonyl (-S(O)₂-);

10

E is N~OH, N~OC(O)CH₃, O, S, with the proviso that when E is bivalent sulfur or oxygen, R_5 and R_6 cannot both be hydrogen;

R_8 is C₁-C₃-alkyl;

15

R_9 is H or C₁-C₃-alkyl; and

R_{10} is H, or C₁-C₃-alkyl.

20

Compounds of the present invention contain one or more asymmetric centers and therefore exist as enantiomers or diastereoisomers. Individual stereoisomers or enantiomers can be obtained from the mixtures by known methods of resolution.

As used herein in formula I:

25

Ar denotes an aromatic or hetero-aromatic group such as pyridine, thiophene, a bicyclic-aromatic group such as naphthalenes, benzofurans, benzothiophenes, a tricyclic-aromatic group such as anthracenes and fluorenes; and halo denotes F, Cl, Br or I.

30

Preferred compounds within the scope of the invention include the cis- and trans-isomers as well as racemates and enantiomers of the following:

(±)-N-[2-(N,N'-dimethylamino)cyclohexyl]-N-methyl-2-(4-trifluoromethylphenyl)acetamide;

35

(±)-N-[2-(N',N'-dimethylamino)cyclohexyl]-N-propyl-2-(3-methoxyphenyl)acetamide;

(±)-N-[2-(N',N'-dimethylamino)cyclohexyl]-N-methyl-2-(4-azidophenyl)acetamide;

(±)-N-[2-(N',N'-dimethylamino)cyclohexyl]-N-methyl-2-(3,4-dichlorophenyl)acetamide;

(±)-N-[2-(N',N'-dimethylamino)cyclohexyl]-N-methyl-2-(4-methoxyphenyl)acetamide;

- (±)-N-[2-(N',N'-dimethylamino)cyclohexyl]-N-methyl-2-(2-naphthyl)acetamide;
 (±)-N-[2-(N-cyclopropyl-N-methylamino)cyclohexyl]-2-(4-azidophenyl)acetamide;
 (±)-N-(2-(3-acetoxy-1-pyrrolidinyl)cyclohexyl)-N-methyl-2-(3,4-dichlorophenyl)acetamide;
 5 (±)-N-[2-(N-pyrrolidinyl)cyclohexyl]-N-methyl-2-(3,4-dichlorophenyl)acetamide;
 (±)-N-[2-(3-hydroxypyrrolidinyl)cyclohexyl]-N-methyl-2-(3,4-dichlorophenyl)acetamide;
 (±)-N-[2-[N'-(3-hydroxy-1-azetidiny)]cyclohexyl]-N-methyl-2-(3,4-dichlorophenyl)acetamide;
 (±)-N-[2-(N',N'-diethylamino)cyclohexyl]-N-methyl-2-(3,4-dichlorophenyl)acetamide;
 10 (±)-N-[2-(N'-pyrrolidinyl)cyclohexyl]-N-methyl-2-(3,4-dichlorophenyl)propionamide;
 (±)-N-[2-(4-methyl-1-piperazinyl)cyclopentyl]-2-(3,4-dichlorophenyl)acetamide;
 (±)-N-[2-(N,N-dimethylamino)cyclohexyl]-2-(3,4-dichlorophenyl)acetamide;
 (±)-3,4-dichloro-N-methyl-N-[8-(1-pyrrolidinyl)-1,4-dioxaspiro[4.5]dec-7-yl]-benzeneacetamide;
 15 (±)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidinyl)-1,4-dioxaspiro[4.5]dec-8-yl]-benzeneacetamide;
 (±)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidinyl)-1,4-dioxaspiro[4.5]dec-6-yl]-benzeneacetamide;
 (±)-4-bromo-N-methyl-N-[7-(1-pyrrolidinyl)-1,4-dioxaspiro[4.5]dec-8-yl]-benzeneacetamide;
 20 (±)-3-fluoro-N-methyl-N-[7-(1-azetidiny)]-1,4-dioxaspiro[4.5]dec-8-yl]benzeneacetamide;
 (±)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidinyl)-1,4-dioxaspiro[4.4]-non-8-yl]-benzeneacetamide;
 (±)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidinyl)-1,4-dioxaspiro[4.6]-undec-8-yl]-benzeneacetamide;
 25 (±)-3,4-dichloro-N-methyl-N-[8-(1-pyrrolidinyl)-1,4-dioxaspiro[4.6]-undec-7-yl]-benzeneacetamide;
 (±)-3,4-dichloro-N-methyl-N-[9-(1-pyrrolidinyl)-1,4-dioxaspiro[4.6]-undec-8-yl]-benzeneacetamide;
 30 (±)-3,4-dichloro-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 (±)-3,4-dichloro-N-[5-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 (±)-3,4-dichloro-N-methyl-N-[4-oxo-2-(1-pyrrolidinyl)cyclohexyl]-benzeneacetamide;
 (±)-4-bromo-N-methyl-N-[2-(N',N'-dimethylamino)-4-oxo-cyclohexyl]benzeneacetamide;
 35 (±)-N-[4-acetyloxy-2-(1-pyrrolidinyl)cyclohexyl]-3,4-dichloro-N-methylbenzeneacetamide;
 (±)-N-[4-acetyloxy-2-aminocyclohexyl]-3,4-difluoro-N-methylbenzeneacetamide;
 (±)-3,4-dichloro-N-[5-(hydroxyimino)-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 40 (±)-3,4-dichloro-N-[4,4-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide
 which can also be named:
 (±)-3,4-dichloro-N-methyl-N-[4-oxo-2-(1-pyrrolidinyl)cyclohexyl]benzeneacetamide,
 dimethyl ketal;
 (±)-3,4-dichloro-N-[5,5-diethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;

- (±)-(1 α , 2 β)-3,4-dichloro-N-[4,4-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl benzeneacetamide;
- (±)-4-trifluoromethyl-N-[4,4-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamides ;
- 5 (±)-3-trifluoromethyl -N-[4,4-diethoxy-2- (1-pyrrolidinyl)-cyclohexyl]-N-methylbenzeneacetamide;
- (±)-3-hydroxy-4-methyl-N-[4,4-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
- (±)-4-methanesulfonyl-N-[4,4-dimethoxy-2-(1-piperidinyl)-cyclohexyl]-N-methylbenzamide;
- 10 (±)-4- acetyloxy-N-[4,4-dimethoxy-2- (1-pyrrolidinyl)-cyclohexyl-N-methylbenzeneacetamide;
- (±)-N-[4,4-bis(methylthio)-2-(1-pyrrolidinyl)cyclohexyl]-3,4-dichloro-N-methylbenzeneacetamide;
- 15 (±)-N-[5,5-bis(ethylthio)-2-(1-pyrrolidinyl)cyclohexyl]-3,4-di-chloro-N-methylbenzeneacetamide;
- (±)-3,4-dichloro-N-[4-methylthio-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
- (±)-3,4-dichloro-N-[5-ethylthio-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
- 20 (±)-3,4-dichloro-N-[6-methylthio-2-(1-pyrrolidinyl)cycloheptyl]-N-methylbenzeneacetamide;
- (±)-3,4-dichloro-N-[4-mercapto-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
- 25 [1R-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-4-benzofuranacetamide;
- [1S-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-4-benzofuranacetamide;
- [1R-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-4-benzofuranacetamide;
- 30 [1S-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-4-benzofuranacetamide;
- [1R-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-benzo[b]thiophene-4-acetamide;
- 35 [1S-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-benzo[b]thiophene-4-acetamide;
- [1R-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-benzo[b]thiophene-4-acetamide;
- [1S-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-benzo[b]thiophene-4-acetamide;
- 40 [1R-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-1-naphthaleneacetamide;
- [1S-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-1-naphthaleneacetamide;

- [1R-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-1-naphthaleneacetamide;
 [1S-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-1-naphthaleneacetamide;
 5 [1R-(1 α ,2 β ,4 β ,5 β)]-3,4-dichloro-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 [1S-(1 α ,2 β ,4 β ,5 β)]-3,4-dichloro-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 10 [1R-(1 α ,2 β ,4 α ,5 α)]-3,4-dichloro-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 [1S-(1 α ,2 β ,4 α ,5 α)]-3,4-dichloro-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 [1R-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-9H-fluorene-9-carboxamide;
 15 [1S-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-9H-fluorene-9-carboxamide;
 [1R-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-9H-fluorene-9-carboxamide;
 20 [1S-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-9H-fluorene-9-carboxamide;
 (\pm)-(1 α ,2 β ,4 β)-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-4-benzofuranacetamide;
 (\pm)-(1 α ,2 β ,4 α)-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-4-benzofuranacetamide;
 (\pm)-(1 α ,2 β ,5 β)-N-methyl-N-[5-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-4-benzofuranacetamide;
 (\pm)-(1 α ,2 β ,5 α)-N-methyl-N-[5-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-4-benzofuranacetamide;
 25 (\pm)-(1 α ,2 β ,4 α)-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-9H-fluorene-9-carboxamide;
 (\pm)-(1 α ,2 β ,5 β)-N-[5-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-9H-fluorene-9-carboxamide;
 (\pm)-N-methyl-2-(1-naphthalenyloxy)-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
 30 (\pm)-N-methyl-2-(2-naphthalenyloxy)-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
 (\pm)-1,2-dihydro-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]-1-acenaphthylencarboxamide, (isomer I, mixture of (1 α , 2 β) and (1 β ,2 α) forms);
 (\pm)-1,2-dihydro-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]-1-acenaphthylencarboxamide, (isomer II, mixture of (1 α , 2 β) and (1 β ,2 α) forms);
 35 (\pm)-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-1,2-dihydro-N-methyl-1-acenaphthylencarboxamide (isomer I, mixture of (1 α ,2 β , 4 β , 5 β) and (1 β , 2 α , 4 α , 5 α) forms);
 (\pm)-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-1,2-dihydro-N-methyl-1-acenaphthylencarboxamide (isomer II, mixture of (1 α ,2 β , 4 β , 5 β) and (1 β , 2 α , 4 α , 5 α) forms);
 40 (\pm)-1,2-dihydro-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-1-acenaphthylencarboxamide (isomers I and II, mixtures of (1 α ,2 β , 4 β) and (1 β , 2 α , 4 α) forms);


- (±)-1,2-dihydro-N-[4-methoxy-2-(1-pyrrolidiny)cyclohexyl]-N-methyl-1-acenaphthylenecarboxamide (isomers I and 11, mixtures of (1β, 2α, 4α) and (1α, 2β, 4β) forms);
- (±)-trans-N-methyl-N-[2-(1-pyrrolidiny)cyclohexyl]-9H-fluorene-9-carboxamide;
- 5 (±)-trans-1,3-dihydro-N-methyl-1-oxo-N-[2-(1-pyrrolidiny)cyclohexyl]-4-isobenzofuranacetamide;
- (±)-(1α, 2β, 4β, 5β)-N-[4,5-dimethoxy-2-(1-pyrrolidiny)cyclohexyl]-1,3-dihydro-N-methyl-1-oxo-4-isobenzofuranacetamide;
- 10 (±)-(5α, 7α, 8β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]benzeneacetamide;
- (±)-(5α, 7α, 8β)-bromo-N-methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]benzeneacetamide;
- (±)-(5α, 7α, 8β)-4-methoxy-N-methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]benzeneacetamide;
- 15 (±)-(5α, 7α, 8β)-N-methyl-2-nitro-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]benzeneacetamide;
- (±)-(5α, 7α, 8β)-N-methyl-3-nitro-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]benzeneacetamide;
- (±)-(5α, 7α, 8β)-N-methyl-4-nitro-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]benzeneacetamide;
- 20 (±)-(5α, 7α, 8β)-N-methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-3-(trifluoromethyl)benzeneacetamide;
- (±)-(5α, 6α, 7β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-6-yl]benzeneacetamide;
- 25 (±)-(5α, 7α, 8β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-thiaspiro[4.5]dec-8-yl]benzeneacetamide;
- (±)-(5α, 7β, 8α)-3,4-dichloro-N-methyl-N-[8-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-7-yl]benzeneacetamide;
- (±)-(5α, 7α, 8β)-3,4-dichloro-1,N-dimethyl-[7-(1-pyrrolidiny)-1-azaspiro[4.5]dec-8-yl]benzeneacetamide;
- 30 (±)-(5α, 7α, 8β)-4-bromo-N-methyl-N-[7-(1-pyrrolidiny)-1-azaspiro[4.5]dec-8-yl]benzamide;
- (±)-(5α, 7α, 8β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-thiaspiro[4.5]dec-8-yl]benzamide;
- 35 (±)-(5α, 7α, 8β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-thiaspiro[4.5]dec-8-yl]benzeneacetamide;
- (±)-(5α, 7α, 8β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-thiaspiro[4.5]dec-8-yl]benzeneacetamide, 1-oxide;
- (±)-(5α, 7α, 8β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-thiaspiro[4.5]dec-8-yl]benzeneacetamide, 1,1-dioxide;
- 40 (±)-(5α, 7α, 8β)-N-methyl-N-[7-(1-pyrrolidiny)-1-azaspiro[4.5]dec-8-yl]4-trifluoromethylbenzeneacetamide;
- (±)-(5α, 7α, 8β)-N-methyl-N-[8-(1-pyrrolidiny)-1-thiaspiro[4.5]dec-7-yl]-3-trifluoromethylbenzeneacetamide;
- 45 [5R-(5α, 7α, 8β)]-N-Methyl-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-1H-indene-3-acetamide;

- [5S-(5 α ,7 α ,8 β)]- N-Methyl-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-1H-indene-3-acetamide;
 [5R-(5 α ,7 β ,8 α)]- N-Methyl-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-1H-indene-3-acetamide;
 [5S-(5 α ,7 β ,8 α)]- N-Methyl-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-1H-indene-3-acetamide;
 [5R-(5 α ,7 α ,8 β)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-1H-indole-3-acetamide;
 5 [5S-(5 α ,7 α ,8 β)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-1H-indole-3-acetamide;
 [5R-(5 α ,7 β ,8 α)]- N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-1H-indole-3-acetamide;
 [5S-(5 α ,7 β ,8 α)]- N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-1H-indole-3-acetamide;
 [5R-(5 α ,7 α ,8 β)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-2-benzofuranacetamide;
 [5S-(5 α ,7 α ,8 β)]-N- Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-2-benzofuranacetamide;
 10 [5R-(5 α ,7 β ,8 α)]-N-Methyl-N-[7-(1 -pyrrolidinyl)-1 -oxaspiro[4.5]dec-8-yl]-2-benzo[b]furanacetamide;
 [5S-(5 α ,7 β ,8 α)]-N-Methyl-N-[7-(1 -pyrrolidinyl)-1 -oxaspiro[4.5]dec-8-yl]-2-benzo[b]furanacetamide;
 [5R-(5 α ,7 α ,8 β)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-3-benzo[b]furanacetamide;
 15 [5S-(5 α ,7 α ,8 β)]-N- Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-3-benzo[b]furanacetamide;
 [5R-(5 α ,7 β ,8 α)]-N-Methyl-N-[7-(1 -pyrrolidinyl)-1 -oxaspiro[4.5]dec-8-yl]-3-benzo[b]furanacetamide;
 20 [5S-(5 α ,7 β ,8 α)]-N-Methyl-N-[7-(1 -pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-3-benzo[b]furanacetamide;
 [5R-(5 α ,7 α ,8 β)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-4-benzo[b]furanacetamide;
 [5S-(5 α ,7 α ,8 β)]-N- Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-4-benzo[b]furanacetamide;
 25 [5R-(5 α ,7 β ,8 α)]-N-Methyl-N-[7-(1 -pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-4-benzo[b]furanacetamide;
 [5S-(5 α ,7 β ,8 α)]-N-Methyl-N-[7-(1 -pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-4-benzo[b]furanacetamide;
 30 [5R-(5 α ,7 α ,8 β)]-N-Methyl-N-7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl-4-benzo[b]thiophene-4-acetamide;
 [5S-(5 α ,7 α ,8 β)]- N-Methyl-N-7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl-4-benzo[b]thiophene-4-acetamide;
 [5R-(5 α ,7 β ,8 α)]- N-Methyl-N-7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl-4-benzo[b]thiophene-4-acetamide;
 35 [5S-(5 α ,7 β ,8 α)]- N-Methyl-N-7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl-4-benzo[b]thiophene-4-acetamide;
 (-)-(5 α ,7 α ,8 β)]-N- Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-4-benzo[b]furanacetamide;
 40 (-)-(5 α ,7 α ,8 β)]-N-7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl-4-benzo[b]thiophene-4-acetamide;
 (\pm)-(5 α ,6 α ,7 β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidinyl)-2-oxaspiro[4.5]dec-6-yl]benzeneacetamide;
 (\pm)-(5 α ,6 α ,7 β)-3,4-dichloro-N-methyl-N-[6-(1-pyrrolidinyl)-2-oxaspiro[4.5]dec-7-yl]benzeneacetamide; and


(±)-(5α,7α,8β)-3,4-dichloro-N-methyl-N-[8-(1-pyrrolidinyl)-2-oxaspiro[4.5]dec-7-yl] benzeneacetamide.

The compounds of Formula II and Formula IIa have the following structure:

5



II




IIa

10

wherein for the enantiomers and racemic mixtures

n is 0 or 1;


A is



15 or, -CH₂CH₂- provided that in Formula II, when n is 1, A may also be -O- or -S-;
B, C and D are independently selected from the group consisting of H, OH, OCOR⁵,
OCH₂CH₂OR⁵, OR⁶, R⁶, CH₂OR⁶, CH₂COR⁷, Cl, F, Br, I, NH₂, NHR⁸, NR⁸R⁹, SH, SR⁶,
CH₂SR⁶ and OC(S)N(CH₃)₂; or

20 two of B, C and D when on adjacent carbon atoms taken together form a fused benzo ring;

X and Y are independently selected from the group consisting of H, OCH₃, Cl, F, Br, I, NO₂, CF₃, CN, SO₂R¹⁰, and SO₂CF₃; or
X and Y taken together with the benzene ring form



5

R and R¹ independently are selected from the group consisting of H, and alkyl of 1 to 3 carbon atoms;

10

R² is H; alkyl of 1 to 6 carbon atoms; CH₂CF₃; alkenylmethyl of 3 to 6 carbon atoms; hydroxyalkenylmethyl of 2 to 5 carbon atoms; cycloalkyl of 3 to 6 carbon atoms; cyclopropylmethyl; cyclobutylmethyl, or phenylalkyl of 7 to 9 carbon atoms; or R² can be taken together with R¹ and the nitrogen to which they are attached to be 1-azetidiny; 1-pyrrolidiny optionally substituted at the 3-position by OH, alkyl of 1 to 3 carbon atoms, alkoxy of 1 to 3 carbon atoms or alkanoyloxy of 1 to 3 carbon atoms; 1-piperazinyl optionally substituted at the 4-position by alkyl of 1 to 3 carbon atoms; 1-morpholino; 2,5-dihydro-1H-pyrrol-1-yl; 3-azabicyclo[3.1.0]hexan-3-yl; or 3-azabicyclo[3.2.0]heptan-3-yl;

20

R³ is H, but if n is 1 and A is CH₂, R³ may also be CH₃, CH₂OH, CHO, or COR¹¹;

R⁴ is H, alkyl of 1 to 6 carbon atoms, -CH₂OH-, CHO, or COR¹²;


R⁵ is alkyl of 1 to 6 carbon atoms, phenyl, or mono-substituted phenyl;

R⁶, R⁸, R⁹, R¹⁰ and R¹³ are independently an alkyl group of 1 to 3 carbon atoms; and

25 R⁷, R¹¹ and R¹² independently are selected from the group consisting of H, OH, OR¹³, NHR¹³, and NR₂¹³; or

a stable N-oxide or a pharmaceutically acceptable salt thereof.

30 Preferred compounds are those of Formula IIb, particularly those having the formula



IIb

wherein

5 n is 1;

A is $-\text{CH}_2-$, $-\text{O}-$, or $-\text{S}-$;

B is OH , OCOR^5 , $\text{OCH}_2\text{CH}_2\text{OR}^5$, OR^6 , CH_2OR^6 , or CH_2COR^7 ;

C is H , OH , or OR^6 ; and

10 R^1 and R^2 independently are selected from H or alkyl of 1 to 3 carbon atoms or are taken together with the nitrogen to which they are attached to form the group 1-azetidiny, 1-pyrrolidiny, 1-(2,5-dihydro-1H-pyrrolyl) or 1-piperidiny.

More preferred are compounds of Formula IIb wherein A is $-\text{CH}_2-$.

15

Specifically preferred compounds include:

(\pm)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-5-methoxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride or the methansulfonic acid salt;

20

(\pm)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;

25

(\pm)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-6-methoxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;


(\pm)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-6-hydroxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;

30

(\pm)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;

- (±)trans-3,4-dichloro-N-methyl-N-[2,3-dihydro-2-(pyrrolidin-1-yl)-1H-inden-1-yl]-benzeneacetamide hydrochloride;
- 5 (±)trans-3,4-dichloro-N-methyl-N-[3,4-dihydro-3-(pyrrolidin-1-yl)-2H-benzopyran-4-yl]-benzeneacetamide hydrochloride;
- (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-5-hydroxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- 10 (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-5-propionyloxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- 15 (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-5-benzoyloxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-6,7-dihydroxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- 20 (±)trans-N-methyl-N-[3,4-dihydro-3-(pyrrolidin-1-yl)-2H-benzopyran-4-yl]-benzeneacetamide hydrochloride;
- (±)trans-3,4-dichloro-N-methyl-N-[3,4-dihydro-8-methoxy-3-(pyrrolidin-1-yl)-2H-benzopyran-4-yl]-benzeneacetamide hydrochloride;
- 25 (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-5-(N,N-dimethylthiocarbamoyloxy)-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- 30 (±)trans-3,4-dichloro-N-methyl-N-[2-(2,5-dihydro-1H-pyrrol-1-yl)-5-methoxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride, and
- (±)trans-3-nitro-N-methyl-N-[2,3-dihydro-2-(pyrrolidin-1-yl)-1H-inden-1-yl]-benzeneacetamide hydrochloride.
- 35

Compounds of Formula III have the following structure:



wherein

n is 0-1; and

5 R is unsubstituted phenyl or phenyl substituted with one to three substituents selected from the group consisting of halogen, C₁₋₆ alkyl, hydroxy, -O-CO-NH₂, -O-CO-NHalkyl, -O-CO-N(alkyl)₂, C₁₋₆ alkoxy, trifluoromethyl, C₁₋₄-alkoxy-C₁₋₄ alkyloxy, carboxy-C₁₋₄ alkyloxy, nitrile, nitro and amino; or mono or dialkyl amino, amide, sulfonamide, carboxamide; or mono or disubstituted carboxamide, ureido; or mono and di-alkylsubstituted ureido; or

10

R represents an alkyl or cycloalkyl group having up to 7 carbon atoms. The cycloalkyl moiety, where present, can be optionally substituted by one or more substituents selected from the group consisting of from hydroxy, amino, amidino, guanidino, aminocarbonyl, carboxy, C₁₋₆ alkoxy, (C₁₋₆ alkoxy)carbonyl, (C₃₋₆ alkenyloxy)carbonyl, (C₃₋₆ alkynyloxy)carbonyl, C₁₋₆ alkanoyloxy, C₁₋₆ alkylsulfide, C₁₋₆ alkylsulfoxide, C₁₋₆ alkylsulfone, C₁₋₆(monoalkylamino)carbonyl, C₁₋₆ acylamino, C₁₋₆ acylmethylamino and C₁₋₆ monoalkylamino; or

15

20 R represents the group -B-R⁷ in which B represents -CH₂-, -CH(CH₃)- or a single bond and R⁷ represents an optionally substituted C₆₋₁₀ carbocyclic aryl group with one to three substituents selected from the group consisting of halogen, C₁₋₆ alkyl, hydroxy, -O-CO-NH₂, -O-CO-NHalkyl, -O-CO-N(alkyl)₂, C₁₋₆ alkoxy, trifluoromethyl, C₁₋₄-alkoxy-C₁₋₄ alkyloxy, carboxy-C₁₋₄ alkyloxy, nitrile, nitro and amino; or mono or dialkyl amino, amide, sulfonamide or carboxamide; mono or disubstituted carboxamide or ureido; and

25

30 R represents the group -D-R⁸ in which D represents a single bond, -CH₂-, -CH(CH₃)-, -CH₂O-, -CH(CH₃)O-, -CH₂S-, -CH(CH₃)S-, -CH₂NH- or -CH(CH₃)NH- and R⁸ represents a 4-6 membered heterocyclic ring containing up to 4 heteroatoms selected from the group consisting of oxygen, sulfur and nitrogen, the heterocyclic ring optionally being substituted on nitrogen or sulfur by oxygen or on nitrogen by hydroxy or C₁₋₃ alkyl and/or the ring optionally being substituted on carbon by one or more substituents selected from

the group consisting of amino, hydroxy, thio (and their tautomers), cyano, halogen, C₁₋₃ alkoxy, C₁₋₃ monoalkylamino, C₁₋₃ acylamino, C₁₋₃ acylmethylamino, and C₁₋₃ alkylthio;

- R¹ and R² are independently selected from the group consisting of H, C₁₋₆ alkyl, C₃₋₅ alkenyl, C₃₋₅ alkynyl, and C₄₋₇ cycloalkylalkyl group; or R² can be taken together with R¹ and the nitrogen to which they are attached to form a heterocyclic ring which may optionally contain a further heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur (i.e. 1-azetidiny; 1-pyrrolidinyl optionally substituted at the 3-position by OH, -CH₂OH, tri(C₁-C₆ alkyl)silyloxy, acyloxy, C₁₋₆ alkyl, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy; 1-piperazinyl optionally substituted at the 4-position by alkyl of 1 to 3 carbon atoms; 1-morpholino; 2,5-dihydro-1H-pyrrol-1-yl; 3-azabicyclo[3.1.0]hexan-3-yl; or 3-azabicyclo[3.2.0]heptan-3-yl);


- R³ represents hydrogen, C₁₋₇ alkyl, -CH₂-phenyl or heterocyclic (the phenyl or heterocyclic substituted with one to three substituents selected from the group consisting of halo, C₁₋₄ alkyl, C₁₋₄ alkoxy and methoxycarbonyl; mono-, di- or tri-halomethyl; cyano; COR⁹; CH=NOR¹⁰; OR¹⁰; SR¹⁰; CH₂CN; CH₂OR¹⁰; CH₂SR¹⁰; CH₂S(O)R¹⁰; CH₂S(O)₂R¹⁰; CH₂N(R¹⁰)R¹¹; CH₂(R¹⁰)R¹¹; CH₂NR¹⁰OH; CH₂N(COR¹⁰)OH; CH₂NR¹⁰COR¹¹; CH₂NR¹⁰S(O)₂R¹¹; or CH₂OCOR¹⁰, wherein R⁹ is hydrogen, hydroxy, amino, NHOH, NHOCH₃, pyridylamino, NHN(CH₃)₂, C₁₋₄ alkoxy, benzyloxy, C₁₋₄ alkylamino, di-C₁₋₄ alkylamino, C₁₋₄ alkyl or C₁₋₄ alkylthio; R¹⁰ and R¹¹ are each hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy or C₇₋₁₁ phenylalkyl), or OR¹², wherein R¹² is hydrogen, C₁₋₄ alkyl or a hydroxy protecting group;

X represents -CO-, or -SO₂-; and

Y represents a single bond (in this case, only one of R⁴-R⁶ is attached), a tetrahedral carbon, -OC-, -SC-, -S(O)C-, -S(O)₂C-, or -CH₂C-; and

- R⁴, R⁵, and R⁶ are independently selected from the group consisting of hydrogen, hydroxy, alkoxy, C₁₋₄ alkylendioxy, C₁₋₈ cyclic and acyclic alkyl; substituted or unsubstituted carbocyclic aromatic or heterocyclic aromatic group (i.e. phenyl, naphthyl, biphenyl, indanyl, 1-tetralone-6-yl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl or benzothienyl, substituted with one to three substituents selected from the group consisting

- of halo, cyano, $-\text{OCONH}_2$, $-\text{OCONHalkyl}$, $-\text{OCON(alkyl)}_2$, $-\text{OCOalkyl}$, $-\text{NHCHO}$, $-\text{NHCOalkyl}$, ureido, $-\text{NHCONHalkyl}$, $-\text{NalkylCONHalkyl}$, $-\text{NHCON(alkyl)}_2$, $-\text{NalkylCON(alkyl)}_2$, $-\text{NHSO}_2\text{alkyl}$, $-\text{COalkyl}$, $-\text{CONH}_2$, $-\text{CONHalkyl}$, $-\text{CON(alkyl)}_2$, $-\text{CH}_2\text{CONH}_2$, $-\text{CH}_2\text{CONHalkyl}$, $-\text{CH}_2\text{CON(alkyl)}_2$, $-\text{OCH}_2\text{CONH}_2$, $-\text{OCH}_2\text{CONHalkyl}$, $-\text{OCH}_2\text{CON(alkyl)}_2$, C_{1-4} alkyl, C_{1-4} alkoxy, amino, hydroxy, nitro, trifluoromethyl, $-\text{SO}_2\text{alkyl}$, $-\text{SOalkyl}$, and mesyl; or R^5 and R^6 can together form the following structure



- wherein R^{13} and R^{14} are independently selected from the group consisting of hydrogen, halogen, hydroxy, alkoxy, mono-, di- or tri-halomethyl, amino, $-\text{NHalkyl}$, $-\text{N(alkyl)}_2$, $-\text{NHCOalkyl}$, ureido, nitro, and methylenedioxy; and


D represents $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{NH}-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}=\text{CH}-$, $-\text{CH}_2\text{NH}-$, or $-\text{CH}_2\text{Nalkyl}-$.

Preferred examples include the following compounds:

- N-methyl-N- $\{[1\text{S}]-1\text{-phenyl-2-}[(3\text{S})-(3\text{-hydroxypyrrolidin-1-yl})]\text{ethyl}\}-2,2\text{-diphenylacetamide hydrochloride}$,
- 15 3,4-dichloro-N-methyl-N- $\{[1\text{S}]-1\text{-phenyl-2-(1-pyrrolidinyl)ethyl}\}$ benzeneacetamide hydrochloride,
- N-methyl-N- $\{[1\text{S}]-1\text{-phenyl-2-}[(3\text{S})-(3\text{-hydroxypyrrolidin-1-yl})]\text{ethyl}\}-2\text{-aminophenylacetamide hydrochloride}$,
- 3,4-dichloro-N-methyl-N- $\{[1\text{S}]-1\text{-isopropyl-2-(1-pyrrolidinyl)ethyl}\}$ benzeneacetamide hydrochloride,
- 20 3,4-dichloro-N-methyl-N- $\{[1\text{S}]-1\text{-(O-acetic acid-3-hydroxyphenyl)-2-(1-pyrrolidinyl)ethyl}\}$ benzeneacetamide hydrochloride, and

N-methyl-N-[(1S)-1-phenyl-2-(1-pyrrolidiny)ethyl]-2,2-diphenylacetamide
hydrochloride

The compounds of Formula IV have the following structure:



IV

wherein:

R_1 and R_2 are the same or different and are hydrogen, C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} cycloalkyl or C_{4-12} cycloalkylalkyl groups, or together form a C_{2-8} branched or linear polymethylene or C_{2-6} alkenylene group, optionally substituted with a hetero-atom; or - NR_1R_2 form a 5-membered (optionally containing an oxygen atom adjacent to the nitrogen) or 6-membered ring, which ring optionally contains one unit of unsaturation and which is unsubstituted or substituted with hydroxy, C_{1-6} acyloxy, oxo, optionally substituted methylene, $-COR_{10}$ (where R_{10} represents C_{1-6} alkyl, $-OR_{11}$ or $-NH R_{11}$ and R_{11} represents hydrogen, C_{1-6} alkyl, aryl, or $Ar(C_{1-6})$ alkyl, or $N=NOR_{12}$ (where R_{12} represents C_{1-6} alkyl);

R_3 is hydrogen, C_{1-6} alkyl or phenyl; or R_3 together with R_1 forms a $-(CH_2)_3-$ or $-(CH_2)_4-$ group;

R_4 is C_{1-6} alkyl, or phenyl;

R_5 is hydrogen, or together with R_4 forms a C_{2-5} linear polymethylene group;

R_6 represents hydroxy, C_{1-6} alkyl, C_{1-6} hydroxyalkyl, C_{1-6} carboxyalkyl, phenyl, oxo, amino, carboxy, amido, $-COR_{13}$, $-CO_2R_{13}$ or $-COCO_2R_{13}$ (where R_{13} represents a hydrogen atom or an unsubstituted or substituted C_{1-10} hydrocarbon moiety); $-NR_xCOR_x$ (where R_x represents C_{1-6} alkyl), optionally substituted ethylene; or R_6 together with the
 5 E atom to which it is attached, forms a 5 or 6-membered ring containing one or more heteroatoms; R_7 is hydrogen, or together with R_6 forms an optionally substituted or unsubstituted single or fused aryl or heterocyclic ring, containing from 5 to 12 ring atoms and comprising up to four heteroatoms in the ring selected from the group consisting of oxygen, nitrogen and sulphur, which may be substituted with hydrogen, C_{1-6} alkyl, -
 10 CH_2OR_{14} , halogen, hydroxy, C_{1-6} alkoxy, C_{1-6} alkoxy carbonyl, thiol, C_{1-6} alkylthio, $-OCOR_{15}$, $-NHCOR_{16}$, $-NHSO_2R_{17}$ or $-CH_2SO_2NR_{18}R_{19}$,
 in which each of R_{14} to R_{19} is independently hydrogen, C_{1-6} alkyl, aryl or aralkyl,

A is aryl or heteroaryl ring, optionally mono or disubstituted with C_{1-6} alkyl, C_{2-6} alkenyl,
 15 C_{1-6} haloalkyl, C_{2-6} haloalkenyl, C_{2-6} haloalkynyl, aryl, aralkyl, hydroxy, C_{1-6} alkoxy, C_{1-6} haloalkoxy, thiol, C_{1-6} alkylthio, C_{1-6} haloalkylthio, halogen, nitro, cyano, carboxy, aryloxy, aralkoxy carbonyl, carbamoyl, sulfonyl or sulfamoyl;

E represents methylene, sulphur, oxygen or an imino ;
 20

R_8 is hydrogen or C_{1-6} alkyl; and

R_9 is hydrogen or together with R_8 may form the group $-(CRaRa)_m-C(=Y)-$ wherein Ra is hydrogen or C_{1-6} alkyl having up to a maximum of 3 alkyl groups;

25 m is 1, 2, or 3; and

Y represents two hydrogens or oxygen.

30 Preferred compounds of Formula IV include the following:

$(\pm)1-(\text{Pyrrolidin-1-yl})\text{methyl-2-(3,4-dichlorophenyl)-acetyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline}$;

(±)8-[(3,4-Dichlorophenyl)acetyl]-7-(1-pyrrolidinylmethyl)-1,4-dioxo-8-aza[4.5]spirodecane;

5 (±)Methyl 4-[3,4-dichlorophenyl]acetyl]-3-(1-pyrrolidinylmethyl)-1-piperazinecarboxylate
1-[(3,4-Dichlorophenyl)acetyl]-2-[(3-oxo-1-pyrrolidinyl)methyl]-piperidine;

10 (±)[S-(RR)]-(-)5-[(3,4-Dichlorophenyl)acetyl]-4,5,6,7-tetrahydro-4[(3-hydroxy-1-pyrrolidinyl)methyl]furo[3,2-c]pyridine;

(±)[S-(RS)]-4-Acetyl-1-[(3,4-dichlorophenyl)acetyl]-2-[(3-hydroxy-1-pyrrolidinyl)methyl]pyridine;

15 (±)2-[(3,4-Dichlorophenyl)acetyl]-1,2,3,4-tetrahydro-1-(1-pyrrolidinyl)methyl)-5-isoquinolinol;.

(±)4-(Pyrrolidin-1-yl)methyl-5-(3,4-dichlorophenyl)acetyl-4,5,6,7-tetrahydrothieno[3,2-c]pyridine;


20 (±)1-[(5,6-Dichloro-3-oxoindan-1-carbonyl)-2-pyrrolidin-1-ylmethyl]piperidine;

(±)2-(3,4-Dichlorophenyl)acetyl-3-(pyridin-1-yl)methyl-decahydroisoquinoline; and

(±)1-(4-Trifluoromethylphenyl)acetyl-2-(3-hydroxypyrrolidin-1-yl)methyl-4,4-dimethyl piperidine.

25

The compounds of Formula V have the following structure:



30 wherein

----- represents a single or double bond;

R¹ represents an alkyl group having 1-5 carbon atoms, a cycloalkylalkyl group having 4-7 carbon atoms, a cycloalkenylalkyl group having 5-7 carbon atoms, an aryl group having 6-12 carbon atoms, an aralkyl group having 7-13 carbon atoms, an alkenyl group having 4-7 carbon atoms, an allyl group, a furan-2-ylalkyl group having 1-5 carbon atoms, or a thiophen-2-ylalkyl group having 1-5 carbon atoms;

R² represents a hydrogen atom, a hydroxy group, a nitro group, an alkanoyloxy group having 1-5 carbon atoms, an alkoxy group having 1-5 carbon atoms, an alkyl group having 1-5 carbon atoms, or -NR⁹R¹⁰ wherein R⁹ represents a hydrogen atom or an alkyl group having 1-5 carbon atoms, and R¹⁰ represents a hydrogen atom; an alkyl group having 1-5 carbon atoms, or -C(=O)R¹¹ wherein R¹¹ represents a hydrogen atom, a phenyl group or an alkyl group having 1-5 carbon atoms;

R³ represents a hydrogen atom, a hydroxy group, an alkanoyloxy group having 1-5 carbon atoms, or an alkoxy group having 1-5 carbon atoms;

A represents -XC(=Y)-, -XC(=Y)Z-, -X-, -XSO₂-, or -OC(OR⁴)R⁴- (where, X, Y and Z each independently represent NR⁴, S or O wherein R⁴ represents a hydrogen atom, a straight-chain or branched chain alkyl group having 1-5 carbon atoms or an aryl group having 6-12 carbon atoms, and wherein R⁴ may be identical or different;

B represents a valence bond, a straight-chain or branched chain alkylene group having 1-14 carbon atoms (which may be substituted with at least one substituent selected from the group consisting of an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, a hydroxy group, fluorine, chlorine, bromine, iodine, an amino group, a nitro group, a cyano group, a trifluoromethyl group and a phenoxy group, and wherein 1 to 3 methylene groups may be replaced with carbonyl groups), an acyclic unsaturated hydrocarbon containing from 1 to 3 double bonds and/or triple bonds and having 2-14 carbon atoms (which may be substituted with at least one substituent group selected from the group consisting of an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, a hydroxy group, fluorine, chlorine, bromine, iodine, an amino group, a nitro group, a cyano group, a trifluoromethyl group and a phenoxy group, and wherein from 1 to 3 methylene groups may be replaced with




carbonyl groups), or a straight-chain or branched chain saturated or unsaturated hydrocarbon group containing from 1 to 5 thioether, ether and/or amino bonds and having 1-14 carbon atoms (wherein hetero atoms are not bonded directly to A, and 1 to 3 methylene groups may be replaced with carbonyl groups);

5








R^5 represents a hydrogen atom or an organic group (which may be substituted with at least one or more substituent groups selected from the group consisting of an alkyl group having 1-5 carbon atoms, an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, a hydroxy group, fluorine, chlorine, bromine, iodine, an amino group, a nitro group, a cyano group, an isothiocyanate group, a trifluoromethyl group and a methylenedioxy group); or

10

R_5 is



15



wherein
Q is N, O or S;

20 T is CH, N, S or O;

l is 0-5;

m and n are ≥ 0

m + n ≤ 5 ;

R⁶ represents a hydrogen atom;

5

R⁷ represents a hydrogen atom, a hydroxy group, an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, or R⁶ and R⁷ together represent -O-, -CH₂- or -S-;

- 10 R⁸ represents a hydrogen atom, an alkyl group having 1-5 carbon atoms, or an alkanoyl group having 1-5 carbon atoms. The general formula (V) includes the (+) form, (-) form and (\pm) form of compounds.

Preferred compounds of Formula V are those wherein:

- 15 R¹ is an alkyl group having 1-5 carbon atoms, a cycloalkylmethyl group having 4-7 carbon atoms, a cycloalkenylmethyl group having 5-7 carbon atoms, a phenylalkyl group having 7-13 carbon atoms, an alkenyl group having 4-7 carbon atoms, an allyl group, a furan-2-yl-alkyl group having 1-5 carbon atoms or a thiophen-2-yl-alkyl group having 1-5 carbon atoms, while particularly preferred compounds are those wherein R¹ is methyl,
20 ethyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclopentenylmethyl, cyclohexenylmethyl, benzyl, phenethyl, trans-2-butenyl, 2-methyl-2-butenyl, allyl, furan-2-yl-methyl or thiophen-2-yl-methyl groups;

- 25 R² is hydrogen, hydroxy, nitro, acetoxy, methoxy, methyl, ethyl, propyl, amino, dimethylamino, acetylamino or benzoylamino groups, while particularly preferred compounds are those wherein R² is hydrogen, hydroxy, nitro, acetoxy, methyl or dimethylamino groups;

R³ is hydrogen, hydroxy, acetoxy or methoxy;


30

A is -NR⁴C(=O)-, -NR⁴C(=S)-, -NR⁴C(=O)O-, -NR⁴C(=O)NR⁴-, -NR⁴C(=S)NR⁴-, -NR⁴C(=O)S-, -OC(=O)-, -OC(=O)O-, -SC(=O)-, -NR⁴-, -O-, -NR⁴SO₂- or -OSO₂-, while

particularly preferred compounds are those wherein R^3 is $-NR^4C(=O)-$, $-NR^4C(=S)-$, $-NR^4C(=O)O-$, $-NR^4C(=O)NR^4-$, $-NR^4C(=S)NR^4-$ or $-NR^4SO_2-$;

R^4 is hydrogen, a straight-chain or branched alkyl group having 1-5 carbon atoms or a phenyl group, while particularly preferred compounds are those wherein R^4 is a straight-chain or branched alkyl group having 1-5 carbon atoms, particularly methyl, ethyl, propyl, isopropyl, butyl and isobutyl groups; or

R^4 is



Formula V-1

wherein

Q is N, O or S;

T is CH, N, S or O;

m and n are ≥ 0 and

$m + n \leq 5$;

B is $-(CH_2)_n-$ ($n = 0-6$), $-(CH_2)_n-C(=O)-$ ($n = 1-4$), $-CH=CH-(CH_2)_n-$ ($n = 0-4$), $-C\equiv C-(CH_2)_n-$ ($n=0-4$), $-CH_2-O-$, $-CH_2-S-$, $-CH_2-O-(CH_2)_2-O-(CH_2)_2-$, $-CH_2-O-CH_2-NH-CH_2-O-CH_2-$ or $-CH_2-O-CH_2-S-CH_2-O-CH_2-$, while particularly preferred compounds are those wherein B is $-(CH_2)_n-$ ($n=0-6$), $-CH=CH(CH_2)_n-$ ($n=0-4$), $-C\equiv C-(CH_2)_n-$ ($n=0-4$), $-CH_2-O-$ or $-CH_2-S-$;

R^5 is hydrogen or an organic group having the basic skeleton indicated in (Formula V-1) (which may be substituted with at least one or more substituent groups selected from the group consisting of an alkyl group having 1-5 carbon atoms, an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, a hydroxy group, fluorine,

chlorine, bromine, an amino group, a nitro group, a cyano group, an isothiocyanate group and a trifluoromethyl group), while particularly preferred compounds are those wherein R^5 is hydrogen, phenyl, 3,4-dichlorophenyl, 4-chlorophenyl, 3-chlorophenyl, 3,4-difluorophenyl, 4-fluorophenyl, 3-fluorophenyl, 2-fluorophenyl, 4-bromophenyl, 3-bromophenyl, 2-bromophenyl, 4-nitrophenyl, 3-nitrophenyl, 2-nitrophenyl, 4-trifluoromethylphenyl, 3-trifluoromethylphenyl, 2-trifluoromethylphenyl, 4-methylphenyl, 3-methylphenyl, 2-methylphenyl, 4-methoxyphenyl, 3-methoxyphenyl, 2-methoxy, 3-furanyl, 2-furanyl, 3-thienyl, 2-thienyl, cyclopentyl or cyclohexyl groups.

10 Examples of preferred compounds include:

17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-3-phenylpropionamido)morphinan;

15 17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-trans-3-(3-furyl)acrylamido)morphinan;

17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-trans-3-cyclohexylacrylamido)morphinan;

20 17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-trans-3-(4-trifluoromethylphenyl)acrylamido)morphinan;

25 17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 α -(N-methyl-trans-3-(3-thiophenyl)acrylamido)morphinan;

17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-trans-3-phenylacrylamido)morphinan;

30 17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-trans-2-hexenamido)morphinan; and

17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-phenylpropiolamido)morphinan.

DETAILED DESCRIPTION OF THE INVENTION

In a composition aspect, the kappa agonist compounds of the present invention for use of their anti-pruritic activity are formulated into parenteral, local and topical formulations.

The compositions are formulated as injectables, as oral and rectal formulations for systemic administration, and for local and topical administration as creams, aqueous or non-aqueous suspension, lotions, emulsions, suspensions or emulsions containing micronized particles, gels, foams aerosols, solids and other suitable vehicles for application to the skin, eyes, lips and mucosa, as suppositories or cream for vaginal administration, and as combinations with bandages, patches, bioadhesives and dressings.

In a method aspect the present invention provides method to treat or prevent pruritus by applying an amount of a compound or composition to a mammal to ameliorate or eliminate pruritus. Thus, the method of the present invention comprises a method of treating pruritus internally or externally present in the mammalian body including: irritation associated with inflammation following local infection, blisters, boils, or acute skin injuries, such as abrasions, burns, superficial cuts, surgical incisions, toothaches, contusions, irritations, inflammatory skin conditions, including but not limited to poison ivy, and allergic rashes and dermatitis and any condition that yields a pruritic state or condition.

FORMULATIONS OF THE PRESENT INVENTION

Effective concentrations of one or more of the compounds of the present invention or pharmaceutically acceptable derivatives thereof are mixed with a suitable pharmaceutical carrier or vehicle for systemic, topical or local administration. Compounds are included in an amount effective for reducing the pruritic state or for which treatment is contemplated. The concentration of active compound in the composition will depend on absorption, inactivation, excretion rates of the active compound, the dosage schedule, and amount administered as well as other factors known to those of skill in the art. For topical and local administration, the dosages are higher, typically at least about 5 to 10 fold, than the amount delivered when administered systemically orally.

The dosage of the compound of Formulas I, II, IIa, IIb, III, IV and V for anti-pruritic purposes is from about 0.001 to about 20 mg/kg body weight of the patient. The compounds of Formulas I, II, IIa, IIb, III, IV and V are conveniently prepared in 5, 10, 25, 50, 75, 100 and 200 mg dosage units for administration for 1 to 4 times a day. Preferred unit dosages are from 0.05 to 10 mg/kg body weight of the patient.

The compounds are administered orally, parenterally, rectally and topically.

Pharmaceutical carriers or vehicles suitable for administration of the compounds and for the methods provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration. In addition, the compounds may be formulated as the sole pharmaceutically active ingredient in the composition or may be combined with other active ingredients.

a) Systemic Formulations

The formulations of the present invention are provided for administration to humans and animals in unit dosage forms, such as tablets, capsules, pills, powders, granules, sterile parenteral solutions or suspensions, and oral solutions or suspensions, and oil-water emulsions containing suitable quantities of a compound of Formulas I, II, IIa, IIb, III, IV and V or pharmacologically acceptable salts thereof.

Pharmaceutical dosage unit forms are prepared to provide from about 0.05 mg to about 500 mg and preferably from about 1.0 to about 200 mg of the essential active ingredient or a combination of essential ingredients per dosage unit form.

Oral pharmaceutical dosage forms are either solid or liquid. The solid dosage forms are tablets, capsules, granules, and bulk powders. Types of oral tablets include

compressed, chewable lozenges and tablets which may be enteric-coated, sugar-coated or film-coated. Capsules may be hard or soft gelatin capsules, while granules and powders may be provided in non-effervescent or effervescent form with the combination of other ingredients known to those skilled in the art.

5 Pharmaceutically acceptable carriers utilized in tablets are binders, lubricants, diluents, disintegrating agents, coloring agents, flavoring agents, and wetting agents. Enteric-coated tablets, due to their enteric-coating, resist the action of stomach acid and dissolve or disintegrate in the neutral or alkaline intestines. Sugar-coated tablets are compressed tablets to which different layers of pharmaceutically acceptable substances
10 have been applied. Film-coated tablets are compressed tablets which have been coated with a water soluble polymer. Multiple compressed tablets are compressed tablets made by more than one compression cycle utilizing the pharmaceutically acceptable substances previously mentioned. Coloring agents may also be used in the above dosage forms. Flavoring and sweetening agents are used in compressed tablets, sugar-coated, multiple
15 compressed and chewable tablets. Flavoring and sweetening agents are especially useful in the formation of chewable tablets and lozenges.

 Examples of binders include glucose solution, acacia mucilage, gelatin solution, sucrose and starch paste. Lubricants include talc, starch, magnesium or calcium stearate, lycopodium and stearic acid. Diluents include, for example, lactose, sucrose, starch,
20 kaolin, salt, mannitol and dicalcium phosphate. Disintegrating agents include corn starch, potato starch, bentonite, methylcellulose, agar and carboxymethylcellulose. Coloring agents include, for example, any of the approved certified water soluble FD and C dyes, mixtures thereof, and water insoluble FD and C dyes suspended on alumina hydrate. Sweetening agents include sucrose, lactose, mannitol and artificial sweetening agents
25 such as sodium cyclamate and saccharin, and any number of spray dried flavors. Flavoring agents include natural flavors extracted from plants such as fruits and synthetic blends of compounds which produce a pleasant sensation. Wetting agents include propylene glycol monostearate, sorbitan monooleate, diethylene glycol monolaurate and polyoxyethylene laural ether. Enteric-coatings include fatty acids, fats, waxes, shellac, ammoniated shellac and cellulose acetate phthalates. Film coatings include
30 hydroxyethylcellulose, sodium carboxymethylcellulose, polyethylene glycol 4000 and cellulose acetate phthalate.

 Liquid oral dosage forms include aqueous solutions, emulsions, suspensions, solutions and/or suspensions reconstituted from non-effervescent granules and
35 effervescent preparations reconstituted from effervescent granules. Aqueous solutions include, for example, elixirs and syrups. Emulsions are either oil-in water or water-in-oil.

Elixirs are clear, sweetened, hydroalcoholic preparations. Pharmaceutically acceptable carriers used in elixirs include solvents. Syrups are concentrated aqueous solutions of a sugar, for example, sucrose, and may contain a preservative. An emulsion is a two-phase system in which one liquid is dispersed in the form of small globules throughout another liquid. Pharmaceutically acceptable carriers used in emulsions are non-aqueous liquids, emulsifying agents and preservatives. Suspensions use pharmaceutically acceptable suspending agents and preservatives. Pharmaceutically acceptable substances used in non-effervescent granules, to be reconstituted into a liquid oral dosage form, include diluents, sweeteners and wetting agents. Pharmaceutically acceptable substance used in effervescent granules, to be reconstituted into a liquid oral dosage form, include organic acids and a source of carbon dioxide. Coloring and flavoring agents are used in all of the above dosage forms.

Solvents include glycerin, sorbitol, ethyl alcohol and syrup. Examples of preservatives include glycerin, methyl and propylparaben, benzoic acid, sodium benzoate and alcohol. Examples of non-aqueous liquids utilized in emulsions include mineral oil and cottonseed oil. Examples of emulsifying agents include gelatin, acacia, tragacanth, bentonite, and surfactants such as polyoxyethylene sorbitan monooleate. Suspending agents include sodium carboxymethylcellulose, pectin, tragacanth, Veegum and acacia. Diluents include lactose and sucrose. Sweetening agents include sucrose, syrups, glycerin and artificial sweetening agents such as sodium cyclamate and saccharin. Wetting agents include propylene glycol monostearate, sorbitan monooleate, diethylene glycol monolaurate and polyoxyethylene lauryl ether. Organic acids include citric and tartaric acid. Sources of carbon dioxide include sodium bicarbonate and sodium carbonate. Coloring agents include any of the approved certified water soluble FD and C dyes, and mixtures thereof. Flavoring agents include natural flavors extracted from plants such fruits, and synthetic blends of compounds which produce a pleasant taste sensation.

Parenteral administration of the formulations of the present invention includes intravenous, subcutaneous and intramuscular administrations.

Preparations for parenteral administration include sterile solutions ready for injection, sterile dry soluble products ready to be combined with a solvent just prior to use, including hypodermic tablets, sterile suspensions ready for injection, sterile dry insoluble products ready to be combined with a vehicle just prior to use and sterile emulsions. The solutions may be either aqueous or nonaqueous.

Pharmaceutically acceptable carriers used in parenteral preparations include aqueous vehicles, nonaqueous vehicles, antimicrobial agents, isotonic agents, buffers,

antioxidants, local anesthetics, suspending and dispersing agents, emulsifying agents, sequestering or chelating agents and other pharmaceutically acceptable substances.

Examples of aqueous vehicles include Sodium Chloride Injection, Ringers Injection, Isotonic Dextrose Injection, Sterile Water Injection, Dextrose and Lactated Ringers Injection. Nonaqueous parenteral vehicles include fixed oils of vegetable origin, cottonseed oil, corn oil, sesame oil and peanut oil. Antimicrobial agents in bacteriostatic or fungistatic concentrations must be added to parenteral preparations packaged in multiple-dose containers which include phenols or cresols, mercurials, benzyl alcohol, chlorobutanol, methyl and propyl p-hydroxybenzoic acid esters, thimerosal, benzalkonium chloride and benzethonium chloride. Isotonic agents include sodium chloride and dextrose. Buffers include phosphate and citrate. Antioxidants include sodium bisulfate. Local anesthetics include procaine hydrochloride. Suspending and dispersing agents include sodium carboxymethylcellulose, hydroxypropyl methylcellulose and polyvinylpyrrolidone. Emulsifying agents include Polysorbate 80 (Tween 80). A sequestering or chelating agent of metal ions include EDTA. Pharmaceutical carriers also include ethyl alcohol, polyethylene glycol and propylene glycol for water miscible vehicles and sodium hydroxide, hydrochloric acid, citric acid or lactic acid for pH adjustment.

The concentration of the pharmaceutically active compound is adjusted so that an injection provides an effective amount to produce the desired pharmacological effect. The exact dose depends on the age, weight and condition of the patient or animal as is known in the art.

The unit-dose parenteral preparations are packaged in an ampoule or a syringe with a needle.

All preparations for parenteral administration must be sterile, as is known and practiced in the art.

Illustratively, intravenous or intraarterial infusion of a sterile aqueous solution containing an active compound is an effective mode of administration. Another embodiment is a sterile aqueous or oily solution or suspension containing an active material injected as necessary to produce the desired pharmacological effect.

Pharmaceutical dosage forms for rectal administration are rectal suppositories, capsules and tablets for systemic effect.

Rectal suppositories are used herein mean solid bodies for insertion into the rectum which melt or soften at body temperature releasing one or more pharmacologically or therapeutically active ingredients.

Pharmaceutically acceptable substances utilized in rectal suppositories are bases or vehicles and agents to raise the melting point.

5 Examples of bases include cocoa butter (theobroma oil), glycerin-gelatin, carbowax, (polyoxyethylene glycol) and appropriate mixtures of mono-, di- and triglycerides of fatty acids. Combinations of the various bases may be used. Agents to raise the melting point of suppositories include spermaceti and wax. Rectal suppositories may be prepared either by the compressed method or by molding. The typical weight of a rectal suppository is about 2 to 3 gm.

10 Tablets and capsules for rectal administration are manufactured using the same pharmaceutically acceptable substance and by the same methods as for formulations for oral administration.

15 The pharmaceutically therapeutically active compounds of Formulas I, II, IIa, IIb, III, IV and V are administered orally, parenterally or rectally in unit-dosage forms or multiple-dosage forms. Unit-dose forms as used herein refers to physically discrete units suitable for human and animal subjects and packaged individually as is known in the art. Each unit-dose contains a predetermined quantity of the therapeutically active compound sufficient to produce the desired therapeutic effect, in association with the required pharmaceutical carrier, vehicle or diluent. Examples of unit-dose forms include ampoules and syringes individually packaged tablet or capsule. Unit-dose forms may be administered in fractions or multiples thereof. A multiple-dose form is a plurality of identical unit-dosage forms packaged in a single container to be administered in segregated unit-dose form. Examples of multiple-dose forms include vials, bottles of tablets or capsules or bottles of pint or gallons. Hence, multiple dose form is a multiple of unit-doses which are not segregated in packaging.

20
25 Compounds of the present invention in formulations may be included with other active compounds to obtain desired combinations of properties. Other active compounds with known pharmacological properties include analgesics such as aspirin, phenacetin acetaminophen, propoxyphene, pentazocine, codeine, meperidine, oxycodone, mefenamic acid, and ibuprofen; muscle relaxants such as methocarbamol, orphenadrine, carisoprodol, meprobamate, chlorphenesin carbamate, diazepam, chlordiazepoxide and chlorzoxazone; analeptics such as caffeine, methylphenidate and pentylenetetrazol; corticosteroids such as methylprednisolone, prednisone, prednisolone and dexamethasone; antihistamines such as chlorpheniramine, cyproheptadine, promethazine and pyrilamine.

35

b) Local and Topical Formulations

Typically a therapeutically effective dosage is formulated to contain a concentration of at least about 0.1% w/w up to about 50% w/w or more, preferably more than 1% w/w of the active compound to the treated tissue. The active ingredient may be administered at once, or may be divided into a number of smaller doses to be administered at intervals of time. It is understood that the precise dosage and duration of treatment is a function of the tissue being treated and may be determined empirically using known testing protocols or by extrapolation from in vivo or in vitro test data. It is to be noted that concentrations and dosage values may also vary with the age of the individual treated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the formulations, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed formulations.

The compound may be suspended in micronized or other suitable form or may be derivatized to produce a more soluble active product or to produce a prodrug. The form of the resulting mixture depends upon a number of factors, including the intended mode of administration and the solubility of the compound in the selected carrier or vehicle. The effective concentration is sufficient for ameliorating the hyperalgesic or other condition and may be empirically determined.

Compounds are typically included at concentrations 0.001% w/w or greater than 1% w/w up to 50% w/w or higher. The concentration is generally greater than the concentration for systemic administration of the compound. Preferable concentrations are in the range of 0.01% w/w to about 25%w/w, more preferably 1% w/w to 25% w/w, yet more preferably greater than about 1% w/w to about 10% w/w, and most preferably greater than 1% w/w up to about 5% w/w. Aqueous suspensions and formulations contain 1% w/w or more.

The resulting mixture may be a solution, suspension, emulsions or the like and are formulated as creams, gels, ointments, emulsions, solutions, elixirs, lotions, suspensions, tinctures, pastes, foams, aerosols, irrigations, sprays, suppositories, bandages, or any other formulations suitable for topical or local administration.

The route of administration herein is topical or local administration, and compositions are formulated in a manner suitable for each route of administration. Preferred modes of administration include topical application to the skin, eyes or mucosa, and local application to the joints, such as by intra-articular injection. Thus, typical

vehicles are those suitable for pharmaceutical or cosmetic application to body surfaces or for local injection.

Pharmaceutical and cosmetic carriers or vehicles suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration. In addition, the compounds may be formulated as the sole pharmaceutically active ingredient in the composition or may be combined with other active ingredients. The active compound is included in the carrier in an amount sufficient to exert a therapeutically useful effect in the absence of serious toxic effects on the treated individual. The effective concentration may be determined empirically by testing the compounds using in vitro and in vivo systems, including the animal models described herein.

For topical administration, the compounds may be formulated in compositions in the form of gels, creams, lotions, solids, solutions or suspensions, or aerosols. Compositions for treating human skin are formulated for topical application with an anti-hyperalgesic effective amount of one or more of the compounds selected as described herein, in an effective concentration range [by weight], between about 0.1% and 80%, preferably 0.1 to 50%, more preferably greater than about 1% up to about 50% or more in a cream, ointment, lotion, gel, solution or solid base or vehicle known in the art to be non-toxic and dermatologically acceptable or suitable for application to the mucosa. Aqueous suspensions are preferably formulated at concentrations greater than about 1% w/w, more preferably 2% w/w.

To formulate a composition, the weight fraction of compound is dissolved, suspended, dispersed or otherwise mixed in a selected vehicle at an effective concentration such that the hyperalgesic condition is relieved or ameliorated. Generally, emollient or lubricating vehicles that help hydrate the skin are more preferred than volatile vehicles, such as ethanol, that dry the skin. Examples of suitable bases or vehicles for preparing compositions for use with human skin are petrolatum, petrolatum plus volatile silicones, lanolin, cold cream [USP], and hydrophilic ointment [USP].

The choice of an acceptable vehicle is largely determined by the mode of application and tissue to be treated. Suitable pharmaceutically and dermatologically acceptable vehicles for topical application include those suited for use include lotions, creams, solutions, gels, tapes and the like. Generally, the vehicle is either organic in nature or an aqueous emulsion and capable of having the selected compound or compounds, which may be micronized, dispersed, suspended or dissolved therein. The vehicle may include pharmaceutically-acceptable emollients, skin penetration enhancers, coloring agents, fragrances, emulsifiers, thickening agents, and solvents.

For local internal administration, such as intra-articular administration, the compounds are preferably formulated as a suspension in an aqueous-based medium, such as isotonicly buffered saline or are combined with a biocompatible support or bioadhesive intended for internal administration.

5

Lotions

The lotions contain an effective concentration of one or more of the compounds. The effective concentration is preferably effective to deliver an anti-hyperalgesic amount, typically at a concentration of between about 0.1 - 50% w/w or more of one or more of the compounds provided herein. The lotions also contain from 1% to 50% w/w, preferably from 3% to 15% w/w of an emollient and the balance water, a suitable buffer, a C₂ or C₃ alcohol, or a mixture of water of the buffer and the alcohol. Any emollients known to those of skill in the art as suitable for application to human skin may be used. These include, but are not limited to, the following:

15 (a) Hydrocarbon oils and waxed, including mineral oil, petrolatum, paraffin, ceresin, ozokerite, microcrystalline wax, polyethylene, and perhydrosqualene.

(b) Silicone oils, including dimethylpolysiloxanes, methylphenylpolysiloxanes, water-soluble and alcohol-soluble silicone-glycol copolymers.

20 (c) Triglyceride fats and oils, including those derived from vegetable, animal and marine sources. Examples include, but are not limited to, castor oil, safflower oil, cotton seed oil, corn oil, olive oil, cod liver oil, almond oil, avocado oil, palm oil, sesame oil and soybean oil.

(d) Acetoglyceride esters, such as acetylated monoglycerides.

(e) Ethoxylated glycerides, such as ethoxylated glyceryl monostearate.

25 (f) Alkyl esters of fatty acids having 10 to 20 carbon atoms. Methyl, isopropyl and butyl esters of fatty acids are useful herein. Examples include, but are not limited to, hexyl laurate, isohexyl laurate, isohexyl palmitate, isopropyl palmitate, isopropyl myristate, decyl oleate, isodecyl oleate, hexadecyl stearate, decyl stearate, isopropyl isostearate diisopropyl adipate, diisohexyl adipate, dihexyldecyl adipate, diisopropyl sebacate, lauryl lactate, myristyl lactate, and cetyl lactate.

(g) Alkenyl esters of fatty acids having 10 to 20 carbon atoms. Examples thereof include, but are not limited to, oleyl myristate, oleyl stearate, and oleyl oleate.

35 (h) Fatty acids having 9 to 22 carbon atoms. Suitable examples include, but are not limited to, pelargonic, lauric, myristic, palmitic, stearic, isostearic, hydroxystearic, oleic, linoleic, ricinoleic, arachidonic, behenic, and erucic acids.

(i) Fatty alcohols having 10 to 20 carbon atoms, such as but not limited to, lauryl, myristyl, cetyl, hexadecyl, stearyl, isostearyl, hydroxystearyl, oleyl, ricinoleyl, behenyl, erucyl, and 2-octyl dodecyl alcohols.

(j) Fatty alcohol ethers, including, but not limited to, ethoxylated fatty alcohols of 10 to 20 carbon atoms, such as, but are not limited to, the lauryl cetyl, stearyl, isostearyl, oleyl, and cholesterol alcohols having attached thereto from 1 to 50 ethylene oxide groups or 1 to 50 propylene oxide groups or mixtures thereof.

(k) Ether-esters, such as fatty acid esters of ethoxylated fatty alcohols.

(l) Lanolin and derivatives, including but not limited to, lanolin, lanolin oil, lanolin wax, lanolin alcohols, lanolin fatty acids, isopropyl lanolate, ethoxylated lanolin, ethoxylated lanolin alcohols, ethoxylated cholesterol, propoxylated lanolin alcohols, acetylated lanolin, acetylated lanolin alcohols, lanolin alcohols linoleate, lanolin alcohols ricinoleate, acetate of lanolin alcohols ricinoleate, acetate of ethoxylated alcohols-esters, hydrogenolysis of lanolin, ethoxylated hydrogenated lanolin, ethoxylated sorbitol lanolin, and liquid and semisolid lanolin absorption bases.

(m) Polyhydric alcohols and polyether derivatives, including, but not limited to, propylene glycol, dipropylene glycol, polypropylene glycol [M.W. 2000-4000], polyoxyethylene polyoxypropylene glycols, polyoxypropylene polyoxyethylene glycols, glycerol, ethoxylated glycerol, propoxylated glycerol, sorbitol, ethoxylated sorbitol, hydroxypropyl sorbitol, polyethylene glycol [M.W. 200-6000], methoxy polyethylene glycols 350, 550, 750, 2000, 5000, poly(ethylene oxide) homopolymers [M.W. 100,000 - 5,000,000], polyalkylene glycols and derivatives, hexylene glycol (2-methyl-2,4-pentanediol), 1,3-butylene glycol, 1,2,6-hexanetriol, ethohexadiol USP (2-ethyl-1,3-hexanediol), C₁₅-C₁₈ vicinal glycol and polyoxypropylene derivatives of trimethylolpropane.

(n) Polyhydric alcohol esters, including, but not limited to, ethylene glycol mono- and di-fatty acid esters, diethylene glycol mono- and di-fatty acid esters, polyethylene glycol [M.W. 200-6000], mono- and di-fatty esters, propylene glycol mono- and di-fatty acid esters, polypropylene glycol 2000 monooleate, polypropylene glycol 2000 monostearate, ethoxylated propylene glycol monostearate, glyceryl mono- and di-fatty acid esters, polyglycerol poly-fatty acid esters, ethoxylated glyceryl monostearate, 1,3-butylene glycol monostearate, 1,3-butylene glycol distearate, polyoxyethylene polyol fatty acid ester, sorbitan fatty acid esters, and polyoxyethylene sorbitan fatty acid esters.

(o) Wax esters, including, but not limited to, beeswax, spermaceti, myristyl myristate, and stearyl stearate and beeswax derivatives, including, but not limited to,

polyoxyethylene sorbitol beeswax, which are reaction products of beeswax with ethoxylated sorbitol of varying ethylene oxide content that form a mixture of ether-esters.

(p) Vegetable waxes, including, but not limited to, carnauba and candelilla waxes.

5 (q) Phospholipids, such as lecithin and derivatives.

(r) Sterols, including, but not limited to, cholesterol and cholesterol fatty acid esters.

(s) Amides, such as fatty acid amides, ethoxylated fatty acid amides, and solid fatty acid alkanolamides.

10 The lotions further preferably contain from 1% w/w to 10% w/w, more preferably from 2% w/w to 5% w/w, of an emulsifier. The emulsifiers can be nonionic, anionic or cationic. Examples of satisfactory nonionic emulsifiers include, but are not limited to, fatty alcohols having 10 to 20 carbon atoms, fatty alcohols having 10 to 20 carbon atoms condensed with 2 to 20 moles of ethylene oxide or propylene oxide, alkyl phenols with 6
15 to 12 carbon atoms in the alkyl chain condensed with 2 to 20 moles of ethylene oxide, mono- and di-fatty acid esters of ethylene oxides mono- and di-fatty acid esters of ethylene glycol wherein the fatty acid moiety contains from 10 to 20 carbon atoms, diethylene glycol, polyethylene glycols of molecular weight 200 to 6000, propylene glycols of molecular weight 200 to 3000, glycerol, sorbitol, sorbitan, polyoxyethylene
20 sorbitol, polyoxyethylene sorbitan and hydrophilic wax esters. Suitable anionic emulsifiers include, but are not limited to, the fatty acid soaps, e.g. sodium, potassium and triethanolamine soaps, wherein the fatty acid moiety contains from 10 to 20 carbon atoms. Other suitable anionic emulsifiers include, but are not limited to, the alkali metal, ammonium or substituted ammonium alkyl sulfates, alkyl arylsulfonates, and alkyl
25 ethoxy ether sulfonates having 10 to 30 carbon atoms in the alkyl moiety. The alkyl ethoxy ether sulfonates contain from 1 to 50 ethylene oxide units. Among satisfactory cationic emulsifiers are quaternary ammonium, morpholinium and pyridinium compounds. Certain of the emollients described in preceding paragraphs also have emulsifying properties. When a lotion is formulated containing such an emollient, an
30 additional emulsifier is not needed, though it can be included in the composition.

The balance of the lotion is water or a C₂ or C₃ alcohol, or a mixture of water and the alcohol. The lotions are formulated by simply admixing all of the components together. Preferably, the compound, is dissolved, suspended or otherwise uniformly dispersed in the mixture.

35 Other conventional components of such lotions may be included. One such additive is a thickening agent at a level from 1% to 10% w/w of the composition.

Examples of suitable thickening agents include, but are not limited to: cross-linked carboxypolymethylene polymers, ethyl cellulose, polyethylene glycols, gum, tragacanth, gum kharaya, xanthan gums and bentonite, hydroxyethyl cellulose, and hydroxypropyl cellulose.

5

Creams

The creams are formulated to contain concentration effective to deliver an anti-pruritic amount of the compound to the treated tissue, typically at between about 0.1%, preferably at greater than 1% up to and greater than 50%, preferably between about 3% and 50%, more preferably between about 5% and 15% of one or more of the compounds provided herein. The creams also contain from 5% to 50%, preferably from 10% to 25%, of an emollient and the remainder is water or other suitable non-toxic carrier, such as an isotonic buffer. The emollients, as described above for the lotions, can also be used in the cream compositions. The cream may also contain a suitable emulsifier, as described above. The emulsifier is included in the composition at a level from 3% to 50%, preferably from 5% to 20%.

10

15

Solutions and suspensions for topical and local administration

The solutions are formulated to contain an amount of one or more compounds effective to deliver an anti-pruritic amount, typically at a concentration of between about 0.1 - 50% w/w, preferably at least more than 1% w/w, more preferably more than 2% w/w of one or more of the compounds provided herein. The balance is water, a suitable organic solvent or other suitable solvent or buffer. Suitable organic materials useful as the solvent or a part of a solvent system are as follows: propylene glycol, polyethylene glycol [M.W. 200-600], polypropylene glycol [M.W. 425-2025], glycerine, sorbitol esters, 1,2,6-hexanetriol, ethanol, isopropanol, diethyl tartrate, butanediol and mixtures thereof. Such solvent systems can also contain water.

20

25

30

35

Solutions or suspensions used for local application can include any of the following components: a sterile diluent, such as water for injection, saline solution, fixed oil, polyethylene glycol, glycerine, propylene glycol or other synthetic solvent; antimicrobial agents, such as benzyl alcohol and methyl parabens; antioxidants, such as ascorbic acid and sodium bisulfite; chelating agents, such as ethylenediaminetetraacetic acid [EDTA]; buffers, such as acetates, citrates and phosphates; and agents for the adjustment of tonicity such as sodium chloride or dextrose. Liquid preparations can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass, plastic or other suitable material. Suitable carriers may include physiological saline or phosphate

buffered saline [PBS], and the suspensions and solutions may contain thickening and solubilizing agents, such as glucose, polyethylene glycol, and polypropylene glycol and mixtures thereof. Liposomal suspensions, may also be suitable as pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled in the art.

These compositions that are formulated as solutions or suspensions may be applied to the skin, or may be formulated as an aerosol or foam and applied to the skin as a spray-on. The aerosol compositions typically contain from 25% to 80% w/w, preferably from 30% to 50% w/w, of a suitable propellant. Examples of such propellants are the chlorinated, fluorinated and chlorofluorinated lower molecular weight hydrocarbons. Nitrous oxide, carbon dioxide, butane, and propane are also used as propellant gases. These propellants are used as understood in the art in a quantity and under a pressure suitable to expel the contents of the container.

Suitably prepared solutions and suspension may also be topically applied to the eyes and mucosa. Solutions, particularly those intended for ophthalmic use, may be formulated as 0.01% - 10% w/w isotonic solutions, pH about 5-7, with appropriate salts, and preferably containing one or more of the compounds herein at a concentration of about 0.1% w/w preferably greater than 1% w/w, up to 50% w/w or more. Suitable ophthalmic solutions are known [see, e.g. U.S. Patent No. 5,116,868, which describes typical compositions of ophthalmic irrigation solutions and solutions for topical application]. Such solutions, which have a pH adjusted to about 7.4, contain, for example, 90-100 mM sodium chloride, 4-6 mM dibasic potassium phosphate, 4-6 mM dibasic sodium phosphate, 8-12 mM sodium citrate, 0.5-1.5 mM magnesium chloride, 1.5-2.5 mM calcium chloride, 15-25 mM sodium acetate, 10-20 mM D.L.-sodium β -hydroxybutyrate and 5-5.5 mM glucose.

The active compounds of the present invention can also be mixed with other active materials, that do not impair the desired action, or with materials that supplement the desired action, including viscoelastic materials, such as hyaluronic acid, which is sold under the trademark HEALON [solution of a high molecular weight (MW of about 3 million) fraction of sodium hyaluronate; manufactured by Pharmacia, Inc. see, e.g., U.S. Patent Nos. 5,292,362, 5,282,851, 5,273,056, 5,229,127, 4,517,295 and 4,328,803], VISCOAT [fluorine-containing (meth) acrylates, such as, 1H, 2H, 2H-heptadecafluorodecylmethacrylate; see, e.g., U.S. Patent Nos. 5,278,126, 5,273,751 and 5,214,080; commercially available from Alcon Surgical, Inc.], ORCOLON [see, e.g., U.S. Patent Nos. 5,273,056; commercially available from Optical Radiation Corporation], methylcellulose, methyl hyaluronate, polyacrylamide and polymethacrylamide [see, e.g.,

U.S. Patent No. 5,273,751]. The viscoelastic materials are present generally in amounts ranging from about 0.5 to 5.0% w/w, preferably 1 to 3% w/w of the conjugate material and serve to coat and protect the treated tissues. The compositions may also include a dye, such as methylene blue or other inert dye, so that the composition can be seen when injected into the eye or contacted with the surgical site during surgery.

Gels

Gel compositions can be formulated by simply admixing a suitable thickening agent to the previously described solution or suspension composition. Examples of suitable thickening agents have been previously described with respect to the lotions.

The gelled compositions contain an effective amount of one or more of an anti-pruritic amount, typically at a concentration of between about 0.1 - 50% w/w or more of one or more of the compounds provided therein; from 5% to 75% w/w, preferably from 10% to 50% w/w, of an organic solvent as previously described; from 0.5% to 20% w/w, preferably from 1% to 10% w/w of the thickening agent; the balance being water or other aqueous carrier.

Solids

Compositions of solid forms may be formulated as stick-type compositions intended for application to the lips or other parts of the body. Such compositions contain an effective amount of one or more of the compounds provided therein. The amount is typically an amount effective to deliver an anti-pruritic amount, typically at a concentration of between about 0.1 - 50% w/w or more of one or more of the compounds provided herein. The solids also contain from about 40% to 98% w/w, preferably from about 50% to 90% w/w, of the previously described emollients. This composition can further contain from 1% to 20% w/w, preferably from 5% to 15% w/w, of a suitable thickening agent, and, if desired or needed, emulsifiers and water or buffers. Thickening agents previously described with respect to lotions are suitably employed in the composition in solid form.

Other ingredients such as preservatives, including methyl-paraben or ethyl-paraben, perfumes, dyes or the like, that are known in the art to provide desirable stability, fragrance or color, or other desirable properties, such as shielding from actinic rays from the sun, to compositions for application to the skin may also be employed in a composition for such topical application.

Combinations and kits

The compounds and compositions containing the compounds may also be coated on bandages, mixed with bioadhesives or included in dressings. Thus, combinations of bandages, bioadhesives, dressings and other such materials and the compositions formulated as described herein are provided. Kits containing these combinations, which may also include compositions containing the above listed agents, are also provided.

Articles of manufacture

The compounds and compositions provided herein may be packaged as articles of manufacture containing packaging material, one or more of the compounds provided herein, which is effective for ameliorating pruritus, within the packaging material, and a label that indicates that the compound, N-oxide, acid, salt or other derivative thereof is used for treating pruritic conditions.

Methods of treatment

Compositions for use with human skin preferably may be applied at least once per day, or if necessary, to achieve the desired result, more often, to the areas of the skin for which treatment is sought. It is understood that the precise treatment regimen depends upon the individual treated and may be ascertained empirically depending upon the formulation, and particularly, the age of the treated individual. Any regimen is acceptable as long as the desired anti-pruritic effects are achieved without substantial deleterious or sustained undesirable side effects.

The methods for treating human skin are practiced by applying to the skin, preferably at least daily, a composition suitable for human skin treatment or treatment of mucosal membranes and other body surface tissues, including the vagina, rectum, mouth, eyes and other such tissues. The compositions may be injected into joints or other inflamed areas.

Compositions may be combined with bandages, bioadhesives and other dressings and applied to the body in combination therewith.

The following examples are included for illustrative purposes only and are not intended to limit the scope of the invention.

Example A - Capsules

	Active Compound	2.5 gm
	Corn starch	23.0 gm
5	Lactose	145.0 gm
	Talc	15.0 gm
	Magnesium stearate	3.0 gm

10 The ingredients were mixed and were encapsulated using techniques practiced in the art.

Example B - Tablet

15	Active Compound	150 gm
	Lactose	125 gm
	Corn starch	50 gm
	Magnesium stearate	2.0 gm
	Liquid Petrolatum	2.0 gm

20

The ingredients were mixed, then put through U.S. Standard Screens to produce fine granules. The granules were compressed into tablets, each tablet containing about 150 mg of an active compound of the present invention.

25

Example C - Syrup

	Active Compound	25 gm
	Lemon Oil	2 ml
	Sucrose	650 gm
30	Citric Acid	4 gm
	Benzoic Acid	3 gm
	Tragacanth	16 gm
	Deionized water q.s.	1000 ml

The ingredients, without the active compound, are dispersed in water to make about 800 to 900 ml of solution. The active compound is then added and the solution is stirred into a syrup. Water is then added to make 1000 ml of the syrup.

5 Example D - Parenteral Solution

	Active Compound	30 gm
	Methylparaben	3 gm
	Propylparaben	1 gm
10	Lidocaine	5 gm
	Deionized water q.s. 1000 ml	

The ingredients are dissolved in water to provide a solution followed by sterilization by filtration.

15

Example E - Rectal Suppository

	Active Compound	80 gm
20	Propylene glycol	95 gm
	Polyethylene glycol 4000	1800 gm

The active compound is added to the propylene glycol and milled until a finely divided uniform mixture is formed. The polyethylene glycol 4000 is melted and the propylene glycol dispersion is added with stirring to obtain a suspension. The suspension is poured into molds, allowed to solidify and removed from the molds for packaging.

25

Example F - Water-washable ointment

30

	Active Compound	1.4 % w/w
	Lanolin alcohol	0.15 w/w
	Emulsifying wax NF	7.5% w/w
	PEG-20 glycerides	5.0% w/w
35	Petrolatum	86.0% w/w

The ingredients are melted together and mixed well until the resulting ointment congeals.

Example G - Oil-in-water cream

	Active Compound	10.0% w/w
	Benzyl alcohol	4.0% w/w
5	Propylene glycol	10.0% w/w
	Polyethylene glycol 400	10.0% w/w
	Petrolatum	20.0% w/w
	Stearyl alcohol	10.0% w/w
	Poloxamer	10.0% w/w
10	Water q.s.	100
	Buffer to pH	7.0% w/w

15 In preparing the oil-in-water cream, water, propylene glycol and polyethylene glycol 400 are heated to about 70 to 80_ C, followed by adding a mixture of petrolatum, stearyl alcohol and poloxamer and the mixture is stirred until homogeneous. The active compound in benzyl alcohol is added and the mixture is homogenized. The pH is then adjusted with a buffer to about 7.0.

20 Example H - Aqueous gel

	Active Compound	10.0% w/w
	Benzyl alcohol	4.0% w/w
	Hydroxyethyl cellulose	3.0% w/w
25	Water q.s.	100
	Buffer to pH	7.0% w/w

30 The aqueous gel is prepared by mixing the active compound, benzyl alcohol and adding the mixture to buffered water. Hydroxyethyl cellulose is then added with stirring until the mixture gels.

TESTING OF THE COMPOSITIONS FOR ANTI-PRURITIC ACTIVITY

35 Testing was performed in a mouse scratch model under blind conditions.

Groups of 8-10 male Swiss albino mice (Hilltop Lab Animals, Inc., Scottsdale, PA), weighing 2.5-2.6 g, were used in the testing. They were housed under controlled

temperature of 23-25°C. Food and water were freely available. Before the experiments, the mice were weighted, put into individual boxes and allowed to acclimate for 30 min.

5 Materials

Vehicle used to dissolve the test compounds: 20% w/w cremafor EL.

To induce scratching Compound 48/80 (Sigma, St. Louis, USA) was used which has been shown to produce an itch sensation in humans (Armstrong et al., J. of Physiol., 120: 326, 1953).

10 The compounds to be tested for anti-pruritic activity were dissolved in the vehicle of 20% w/w cremafor EL.

Method

15 100 µl of the vehicle (3-5 doses, n=8-10) was injected s.c. into the back of the neck of mice 20 min. before challenging them with 100 µl of Compound 48/80 (2 mg/ml; 50 µg) injected s.c. into the back of the neck. One minute later the mice were observed for 30 min. and the number of hindleg scratching movements directed to the neck was counted.



20 The vehicle-injected mice scratched 79 ± 16 times in the 30 min after the standard challenge with Compound 48/80.

To each mouse of a group of 8-10 mice previously subjected to the standard challenge various doses of the compounds, to be tested for anti-pruritic activity, were administered s.c. into the back of the neck. One minute later the mice were observed for
25 30 min and the number of hindleg scratching movements directed to the neck was counted.

For each group of 8-10 mice, the mean values for scratching were normalized to relative % antagonism of scratching and then plotted vs. dose of test compounds. Interval estimates of mean A_{50} were determined by nonlinear regression analysis (Kaleida Graph)
30 and mean % inhibition of scratching was calculated.

Representative results are shown in Table I.

TABLE I

Compound	mg/kg s.c.	% Inhibition
	2.5	30
	5.0	57
	10.0	75
	30.0	92
	2.5	24
	5.0	72
	10.0	85


5 Other compounds tested have shown anti-pruritic, dose-responsive activity in the range of from about 15 to about 95% based on doses of from about 0.5 to 10.0 mg/kg, s.c.

It should be understood by those skilled in the art that, while the invention has been described and illustrated above in connection with certain specific embodiments, many variations and modifications may be employed without departing from the scope of
 10 the invention.

WHAT IS CLAIMED IS:

1. An anti-pruritic composition comprising a compound of formula I or a pharmaceutically acceptable salt thereof

5

**I**

wherein

the wavy line bond (~) between the nitrogen in the 2-position carbon indicates the bond can be either cis- or trans with respect to each substituent on the cyclohexyl ring;

A is a single chemical bond (-), $-(CH_2)_q$, $CH(CH_3)-$ or $-X(CH_2)_n$

where q is 1 to 4,
n is 1-4 and
x is O or S;

Ar is an aromatic, hetero-aromatic, bicyclic-aromatic, tricyclic-aromatic group or diphenyl methyl each of which may be unsubstituted or substituted with a member selected from the group consisting of H, halo, trifluoromethyl, nitro, C_1 - C_3 -alkoxy, hydroxy, azido, C_1 - C_3 -alkyl, methanesulfonyl, cyano, amino, C_1 - C_3 -alkoxycarbonyl, C_1 - C_3 -alkanoyloxy, and C_1 - C_3 -carboxacylamino of the formula $-NHC(O)R_7$

where R_7 is H, C_1 - C_2 -alkyl, and aromatic or hetero-aromatic group;


R_1 and R_2 are independently H, C_1 - C_3 -alkyl or allyl;

R_1 and R_2 , taken together with the nitrogen to which they are bonded, complete a ring selected from the group consisting of azetidiny, pyrrolidinyl, 3-hydroxypyrrolidinyl, 3-fluoropyrrolidinyl, morpholinyl, piperidinyl, and 3,4-dehydropiperidinyl;

R_3 , R_4 , R_5 , R_6 are independently H, hydroxy, OR_8 or $OC(=O)R_9$;

R_5 and R_6 taken together may form the group $-E-CH_2-CH_2-E-$;

R_5 and R_6 taken together may form a ring



where

5 Z is selected from the group consisting of oxygen (-O-), NR_{10} , sulfur (-S-), sulfinyl (-S(O)-), and sulfonyl (-S(O)₂-);

E is N~OH, N~OC(O)CH₃, O, S, with the proviso that when E is bivalent sulfur or oxygen, R₅ and R₆ cannot both be hydrogen;

10 R₈ is C₁-C₃-alkyl;

R₉ is H or C₁-C₃-alkyl;

R₁₀ is H, or C₁-C₃-alkyl,

15

in a pharmaceutically acceptable carrier.

2. The anti-pruritic pharmaceutical composition of claim 1 wherein Ar is pyridine, thiophene, naphthalene, benzofuran, benzothiophene anthracene or fluorene; and halo is
20 F, Cl, Br or I.

3. The anti-pruritic pharmaceutical composition of claim 1 wherein said compound is selected from the group consisting of:

25

(±)-N-[2-(N,N'-dimethylamino)cyclohexyl]-N-methyl-2-(4-trifluoromethylphenyl)acetamide;

(±)-N-[2-(N,N'-dimethylamino)cyclohexyl]-N-propyl-2-(3-methoxyphenyl)acetamide;

(±)-N-[2-(N,N'-dimethylamino)cyclohexyl]-N-methyl-2-(4-azidophenyl)acetamide;

30 (±)-N-[2-(N,N'-dimethylamino)cyclohexyl]-N-methyl-2-(3,4-dichlorophenyl)acetamide;

(±)-N-[2-(N,N'-dimethylamino)cyclohexyl]-N-methyl-2-(4-methoxyphenyl)acetamide;

(±)-N-[2-(N,N'-dimethylamino)cyclohexyl]-N-methyl-2-(2-naphthyl)acetamide;

(±)-N-[2-(N-cyclopropyl-N-methylamino)cyclohexyl]-2-(4-azidophenyl)acetamide;

35 (±)-N-(2-(3-acetoxy-1-pyrrolidinyl)cyclohexyl)-N-methyl-2-(3,4-dichlorophenyl)acetamide;

(±)-N-[2-(N-pyrrolidinyl)cyclohexyl]-N-methyl-2-(3,4-dichlorophenyl)acetamide;

(±)-N-[2-(3-hydroxypyrrolidinyl)cyclohexyl]-N-methyl-2-(3,4-dichlorophenyl)acetamide;

(±)-N-[2-[N'-(3-hydroxy-1-azetidyl)cyclohexyl]methyl-2-(3,4-dichlorophenyl)acetamide;

- (±)-N-[2-(N',N'-diethylamino)cyclohexyl]-N-methyl-2-(3,4-dichlorophenyl)acetamide;
 (±)-N-[2-(N'-pyrrolidinyl)cyclohexyl]-N-methyl-2-(3,4-dichlorophenyl)propionamide;
 (±)-N-[2-(4-methyl-1-piperazinyl)cyclopentyl]-2-(3,4-dichlorophenyl)acetamide;
 (±)-N-[2-(N,N-dimethylamino)cyclohexyl]-2-(3,4-dichlorophenyl)acetamide;
 5 (±)-3,4-dichloro-N-methyl-N-[8-(1-pyrrolidinyl)-1,4-dioxaspiro[4.5]dec-7-yl]-benzeneacetamide;
 (±)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidinyl)-1,4-dioxaspiro[4.5]dec-8-yl]-benzeneacetamide;
 (±)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidinyl)-1,4-dioxaspiro[4.5]dec-6-yl]-benzeneacetamide;
 10 (±)-4-bromo-N-methyl-N-[7-(1-pyrrolidinyl)-1,4-dioxaspiro[4.5]dec-8-yl]-benzeneacetamide;
 (±)-3-fluoro-N-methyl-N-[7-(1-azetidyl)-1,4-dioxaspiro[4.5]dec-8-yl]benzeneacetamide;
 (±)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidinyl)-1,4-dioxaspiro[4.4]-non-8-yl]-benzeneacetamide;
 15 (±)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidinyl)-1,4-dioxaspiro[4.6]-undec-8-yl]-benzeneacetamide;
 (±)-3,4-dichloro-N-methyl-N-[8-(1-pyrrolidinyl)-1,4-dioxaspiro[4.6]-undec-7-yl]-benzeneacetamide;
 20 (±)-3,4-dichloro-N-methyl-N-[9-(1-pyrrolidinyl)-1,4-dioxaspiro[4.6]-undec-8-yl]-benzeneacetamide;
 (±)-3,4-dichloro-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 (±)-3,4-dichloro-N-[5-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 25 (±)-3,4-dichloro-N-methyl-N-[4-oxo-2-(1-pyrrolidinyl)cyclohexyl]-benzeneacetamide;
 (±)-4-bromo-N-methyl-N-[2-(N',N'-dimethylamino)-4-oxo-cyclohexyl]benzeneacetamide;
 (±)-N-[4-acetyloxy-2-(1-pyrrolidinyl)cyclohexyl]-3,4-dichloro-N-methylbenzeneacetamide;
 (±)-N-[4-acetyloxy-2-aminocyclohexyl]-3,4-difluoro-N-methylbenzeneacetamide;
 30 (±)-3,4-dichloro-N-[5-(hydroxyimino)-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 (±)-3,4-dichloro-N-[4,4-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide
 which can also be named:
 (±)-3,4-dichloro-N-methyl-N-[4-oxo-2-(1-pyrrolidinyl)cyclohexyl]benzeneacetamide,
 35 dimethyl ketal;
 (±)-3,4-dichloro-N-[5,5-diethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 (±)-(1 α , 2 β)-3,4-dichloro-N-[4,4-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 (±)-4-trifluoromethyl-N-[4,4-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamides ;
 40 (±)-3-trifluoromethyl-N-[4,4-diethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 (±)-3-hydroxy-4-methyl-N-[4,4-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;

- (±)-4-methanesulfonyl-N-[4,4-dimethoxy-2-(1-piperidinyl)-cyclohexyl]-N-methylbenzamide;
- (±)-4-acetyloxy-N-[4,4-dimethoxy-2-(1-pyrrolidinyl)-cyclohexyl]-N-methylbenzeneacetamide;
- 5 (±)-N-[4,4-bis(methylthio)-2-(1-pyrrolidinyl)cyclohexyl]-3,4-dichloro-N-methylbenzeneacetamide;
- (±)-N-[5,5-bis(ethylthio)-2-(1-pyrrolidinyl)cyclohexyl]-3,4-dichloro-N-methylbenzeneacetamide;
- 10 (±)-3,4-dichloro-N-[4-methylthio-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
- (±)-3,4-dichloro-N-[5-ethylthio-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
- (±)-3,4-dichloro-N-[6-methylthio-2-(1-pyrrolidinyl)cycloheptyl]-N-methylbenzeneacetamide;
- 15 (±)-3,4-dichloro-N-[4-mercapto-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
- [1R-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-4-benzofuranacetamide;
- [1S-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-4-benzofuranacetamide;
- 20 [1R-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-4-benzofuranacetamide;
- [1S-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-4-benzofuranacetamide;
- 25 [1R-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzo[b]thiophene-4-acetamide;
- [1S-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzo[b]thiophene-4-acetamide;
- [1R-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzo[b]thiophene-4-acetamide;
- 30 [1S-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzo[b]thiophene-4-acetamide;
- [1R-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-1-naphthaleneacetamide;
- 35 [1S-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-1-naphthaleneacetamide;
- [1R-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-1-naphthaleneacetamide;
- [1S-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-1-naphthaleneacetamide;
- 40 [1R-(1 α ,2 β ,4 β ,5 β)]-3,4-dichloro-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
- [1S-(1 α ,2 β ,4 β ,5 β)]-3,4-dichloro-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;


- [1R-(1 α ,2 β ,4 α ,5 α)]-3,4-dichloro-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 [1S-(1 α ,2 β ,4 α ,5 α)]-3,4-dichloro-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methylbenzeneacetamide;
 5 [1R-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-9H-fluorene-9-carboxamide;
 [1S-(1 α ,2 β ,4 β ,5 β)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-9H-fluorene-9-carboxamide;
 10 [1R-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-9H-fluorene-9-carboxamide;
 [1S-(1 α ,2 β ,4 α ,5 α)]-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-9H-fluorene-9-carboxamide;
 (±)-(1 α ,2 β ,4 β)-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-4-benzofuranacetamide;
 (±)-(1 α ,2 β ,4 α)-N-methyl-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-4-benzofuranacetamide;
 15 (±)-(1 α ,2 β ,5 β)-N-methyl-N-[5-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-4-benzofuranacetamide;
 (±)-(1 α ,2 β ,5 α)-N-methyl-N-[5-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-4-benzofuranacetamide;
 (±)-(1 α ,2 β ,4 α)-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-9H-fluorene-9-carboxamide;
 (±)-(1 α ,2 β ,5 β)-N-[5-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-9H-fluorene-9-carboxamide;
 20 (±)-N-methyl-2-(1-naphthalenyloxy)-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
 (±)-N-methyl-2-(2-naphthalenyloxy)-N-[2-(1-pyrrolidinyl)cyclohexyl]acetamide;
 (±)-1,2-dihydro-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]-1-acenaphthylencarboxamide, (isomer I, mixture of (1 α , 2 β) and (1 β ,2 α) forms);
 25 (±)-1,2-dihydro-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]-1-acenaphthylencarboxamide, (isomer II, mixture of (1 α , 2 β) and (1 β ,2 α) forms);
 (±)-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-1,2-dihydro-N-methyl-1-acenaphthylencarboxamide (isomer I, mixture of (1 α ,2 β , 4 β , 5 β) and (1 β , 2 α , 4 α , 5 α) forms);
 30 (±)-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-1,2-dihydro-N-methyl-1-acenaphthylencarboxamide (isomer II, mixture of (1 α ,2 β , 4 β , 5 β) and (1 β , 2 α , 4 α , 5 α) forms);
 (±)-1,2-dihydro-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-1-acenaphthylencarboxamide (isomers I and II, mixtures of (1 α ,2 β , 4 β) and (1 β , 2 α , 4 α) forms);
 35 (±)-1,2-dihydro-N-[4-methoxy-2-(1-pyrrolidinyl)cyclohexyl]-N-methyl-1-acenaphthylencarboxamide (isomers I and II, mixtures of (1 β , 2 α , 4 α) and (1 α ,2 β , 4 β) forms);
 (±)-trans-N-methyl-N-[2-(1-pyrrolidinyl)cyclohexyl]-9H-fluorene-9-carboxamide;
 40 (±)-trans-1,3-dihydro-N-methyl-1-oxo-N-[2-(1pyrrolidinyl)cyclohexyl]-4-isobenzofuranacetamide;
 (±)-(1 α ,2 β , 4 β , 5 β)-N-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]-1,3-dihydro-N-methyl-1-oxo-4-isobenzofuranacetamide;
 (±)-(5 α ,7 α , 8 β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]benzeneacetamide;
 45

- (±)-(5α,7α, 8β)-bromo-N-methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl] benzeneacetamide;
- (±)-(5α,7α, 8β)-4-methoxy-N-methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl] benzeneacetamide;
- 5 (±)-(5α,7α, 8β)-N-methyl-2-nitro-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl] benzeneacetamide;
- (±)-(5α,7α, 8β)-N-methyl-3-nitro-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl] benzeneacetamide;
- (±)-(5α,7α, 8β)-N-methyl-4-nitro-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl] benzeneacetamide;
- 10 (±)-(5α,7α, 8β)-N-methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-3-(trifluoromethyl)benzeneacetamide;
- (±)-(5α,6α, 7β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-6-yl] benzeneacetamide;
- 15 (±)-(5α,7α, 8β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-thiaspiro[4.5]dec-8-yl] benzeneacetamide;
- (±)-(5α,7β, 8α)-3,4-dichloro-N-methyl-N-[8-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-7-yl] benzeneacetamide;
- (±)-(5α,7α, 8β)-3,4-dichloro-1,N-dimethyl-[7-(1-pyrrolidiny)-1-azaspiro[4.5]dec-8-yl] benzeneacetamide;
- 20 (±)-(5α,7α, 8β)-4-bromo-N-methyl-N-[7-(1-pyrrolidiny)-1-azaspiro[4.5]dec-8-yl] benzamide;
- (±)-(5α,7α, 8β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-thiaspiro[4.5]dec-8-yl] benzamide;
- 25 (±)-(5α,7α, 8β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-thiaspiro[4.5]dec-8-yl] benzeneacetamide;
- (±)-(5α,7α, 8β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-thiaspiro[4.5]dec-8-yl] benzeneacetamide, 1-oxide;
- (±)-(5α,7α, 8β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidiny)-1-thiaspiro[4.5]dec-8-yl] benzeneacetamide, 1,1-dioxide;
- 30 (±)-(5α,7α, 8β)-N-methyl-N-[7-(1-pyrrolidiny)-1-azaspiro[4.5]dec-8-yl]4-trifluoromethylbenzeneacetamide;
- (±)-(5α,7α, 8β)-N-methyl-N-[8-(1-pyrrolidiny)-1-thiaspiro[4.5]dec-7-yl]-3-trifluoromethylbenzeneacetamide;
- 35 [5R-(5α,7α,8β)]-N-Methyl-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-1H-indene-3-acetamide;
- [5S-(5α,7α,8β)]- N-Methyl-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-1H-indene-3-acetamide;
- [5R-(5α,7β,8α)]- N-Methyl-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-1H-indene-3-acetamide;
- [5S-(5α,7β,8α)]- N-Methyl-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-1H-indene-3-acetamide;
- [5R-(5α,7α,8β)]-N-Methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-1H-indole-3-acetamide;
- 40 [5S-(5α,7α,8β)]-N-Methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-1H-indole-3-acetamide;
- [5R-(5α,7β,8α)]- N-Methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-1H-indole-3-acetamide;
- [5S-(5α,7β,8α)]- N-Methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-1H-indole-3-acetamide;
- [5R-(5α,7α,8β)]-N-Methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-2-benzofuranacetamide;
- [5S-(5α,7α,8β)]-N- Methyl-N-[7-(1-pyrrolidiny)-1-oxaspiro[4.5]dec-8-yl]-2-benzofuranacetamide;


- [5R-(5 α ,7 β ,8 α)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-2-benzo[b]furanacetamide;
 [5S-(5 α ,7 β ,8 α)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-2-benzo[b]furanacetamide;
 5 [5R-(5 α ,7 α ,8 β)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-3-benzo[b]furanacetamide;
 [5S-(5 α ,7 α ,8 β)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-3-benzo[b]furanacetamide;
 10 [5R-(5 α ,7 β ,8 α)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-3-benzo[b]furanacetamide;
 [5S-(5 α ,7 β ,8 α)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-3-benzo[b]furanacetamide;
 [5R-(5 α ,7 α ,8 β)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-4-benzo[b]furanacetamide;
 15 [5S-(5 α ,7 α ,8 β)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-4-benzo[b]furanacetamide;
 [5R-(5 α ,7 β ,8 α)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-4-benzo[b]furanacetamide;
 [5S-(5 α ,7 β ,8 α)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-4-benzo[b]furanacetamide;
 20 [5R-(5 α ,7 α ,8 β)]-N-Methyl-N-7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl-4-benzo[b]thiophene-4-acetamide;
 [5S-(5 α ,7 α ,8 β)]-N-Methyl-N-7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl-4-benzo[b]thiophene-4-acetamide;
 25 [5R-(5 α ,7 β ,8 α)]-N-Methyl-N-7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl-4-benzo[b]thiophene-4-acetamide;
 [5S-(5 α ,7 β ,8 α)]-N-Methyl-N-7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl-4-benzo[b]thiophene-4-acetamide;
 (-)-(5 α ,7 α ,8 β)]-N-Methyl-N-[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]-4-benzo[b]furanacetamide;
 30 (-)-(5 α ,7 α ,8 β)]-N-7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl-4-benzo[b]thiophene-4-acetamide;
 (\pm)-(5 α ,6 α ,7 β)-3,4-dichloro-N-methyl-N-[7-(1-pyrrolidinyl)-2-oxaspiro[4.5]dec-6-yl]benzeneacetamide;
 (\pm)-(5 α ,6 α ,7 β)-3,4-dichloro-N-methyl-N-[6-(1-pyrrolidinyl)-2-oxaspiro[4.5]dec-7-yl]benzeneacetamide; and
 35 (\pm)-(5 α ,7 α ,8 β)-3,4-dichloro-N-methyl-N-[8-(1-pyrrolidinyl)-2-oxaspiro[4.5]dec-7-yl]benzeneacetamide.
- 40 4. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of a composition of claim 1.

5. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of a composition of claim 2.
- 5 6. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of a composition of claim 3.
- 10 7. The method of claim 4 wherein said administration is topical administration.
8. The method of claim 4 wherein said administration is parenteral administration.
9. The method of claim 4 wherein said administration is oral administration.
10. The method of claim 4 wherein said administration is rectal administration.
11. The method of claim 5 wherein said administration is topical administration.
- 15 12. The method of claim 5 wherein said administration is parenteral administration.
13. The method of claim 5 wherein said administration is oral administration.
14. The method of claim 5 wherein said administration is rectal administration.
15. The method of claim 6 wherein said administration is topical administration.
16. The method of claim 6 wherein said administration is parenteral administration.
- 20 17. The method of claim 6 wherein said administration is oral administration.
18. The method of claim 6 wherein said administration is rectal administration.
19. An anti-pruritic pharmaceutical composition comprising a compound of formulae II or IIa or a stable N-oxide pharmaceutically acceptable salt thereof

25



II




IIa

- 5 wherein for the enantiomers and racemic mixtures

n is 0 or 1;

A is




or, $-\text{CH}_2\text{CH}_2-$ provided that in Formula II, when n is 1, A may also be $-\text{O}-$ or $-\text{S}-$;

- 10 B, C and D are independently selected from the group consisting of H, OH, OCOR^5 , $\text{OCH}_2\text{CH}_2\text{OR}^5$, OR^6 , R^6 , CH_2OR^6 , CH_2COR^7 , Cl, F, Br, I, NH_2 , NHR^8 , NR^8R^9 , SH, SR^6 , CH_2SR^6 and $\text{OC}(\text{S})\text{N}(\text{CH}_3)_2$; or

- 15 two of B, C and D when on adjacent carbon atoms taken together form a fused benzo ring;

X and Y are independently selected from the group consisting of H, OCH_3 , Cl, F, Br, I, NO_2 , CF_3 , CN, SO_2R^{10} , and SO_2CF_3 ; or

X and Y taken together with the benzene ring form




R and R¹ independently are selected from the group consisting of H, and alkyl of 1 to 3 carbon atoms;

- 5 R² is H; alkyl of 1 to 6 carbon atoms; CH₂CF₃; alkenylmethyl of 3 to 6 carbon atoms; hydroxyalkenylmethyl of 2 to 5 carbon atoms; cycloalkyl of 3 to 6 carbon atoms; cyclopropylmethyl; cyclobutylmethyl, or phenylalkyl of 7 to 9 carbon atoms; or R² can be taken together with R¹ and the nitrogen to which they are attached to form 1-azetidiny, 1-pyrrolidiny optionally substituted at the 3-position by OH, alkyl of 1 to 3 carbon atoms,
- 10 alkoxy of 1 to 3 carbon atoms or alkanoyloxy of 1 to 3 carbon atoms; 1-piperazinyl optionally substituted at the 4-position by alkyl of 1 to 3 carbon atoms; 1-morpholino; 2,5-dihydro-1H-pyrrol-1-yl; 3-azabicyclo[3.1.0]hexan-3-yl; or 3-azabicyclo[3.2.0]heptan-3-yl;
- 15 R³ is H, but if n is 1 and A is CH₂, R³ may also be CH₃, CH₂OH, CHO, or COR¹¹;
R⁴ is H, alkyl of 1 to 6 carbon atoms, -CH₂OH-, CHO, or COR¹²;
R⁵ is alkyl of 1 to 6 carbon atoms, phenyl, or mono-substituted phenyl;
R⁶, R⁸, R⁹, R¹⁰ and R¹³ are independently an alkyl group of 1 to 3 carbon atoms; and
R⁷, R¹¹ and R¹² independently are selected from the group consisting of H, OH, OR¹³,
- 20 NHR¹³, and NR₂¹³;

in a pharmaceutically acceptable vehicle.

20. An anti-pruritic pharmaceutical composition comprising a compound of formulae
- 25 IIb or a stable N-oxide pharmaceutically acceptable salt thereof



IIb

wherein
n is 1;


A is $-\text{CH}_2-$, $-\text{O}-$, or $-\text{S}-$;

5 B is OH, OCOR^5 , $\text{OCH}_2\text{CH}_2\text{OR}^5$, OR^6 , CH_2OR^6 , or CH_2COR^7 ;

C is H, OH, or OR^6 ;

R^1 and R^2 independently are selected from H or alkyl of 1 to 3 carbon atoms or are taken
together with the nitrogen to which they are attached to form the group 1-azetidiny, 1-
10 pyrrolidiny, 1-(2,5-dihydro-1H-pyrrolyl) or 1-piperidiny;

X and Y taken together with the benzene ring form



15

R is H, and $\text{C}_1\text{-C}_3$ alkyl;

R^3 is H; and

R^5 , R^6 and R^7 are independently $\text{C}_1\text{-C}_3$ alkyl,

20 in a pharmaceutically acceptable carrier.

21. The anti-pruritic pharmaceutical composition of claim 19 wherein said compound
is selected from the group consisting of:

25

(\pm)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-5-methoxy-1,2,3,4-
tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride or the methansulfonic acid salt;

- (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- 5 (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-6-methoxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-6-hydroxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- 10 (±)-trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- 15 (±)trans-3,4-dichloro-N-methyl-N-[2,3-dihydro-2-(pyrrolidin-1-yl)-1H-inden-1-yl]-benzeneacetamide hydrochloride;
- (±)trans-3,4-dichloro-N-methyl-N-[3,4-dihydro-3-(pyrrolidin-1-yl)-2H-benzopyran-4-yl]-benzeneacetamide hydrochloride;
- 20 (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-5-hydroxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-5-propionyloxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- 25 (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-5-benzoyloxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- 30 (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-6,7-dihydroxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- (±)trans-N-methyl-N-[3,4-dihydro-3-(pyrrolidin-1-yl)-2H-benzopyran-4-yl]-benzeneacetamide hydrochloride;
- 35 (±)trans-3,4-dichloro-N-methyl-N-[3,4-dihydro-8-methoxy-3-(pyrrolidin-1-yl)-2H-benzopyran-4-yl]-benzeneacetamide hydrochloride;
- (±)trans-3,4-dichloro-N-methyl-N-[2-(pyrrolidin-1-yl)-5-(N,N-dimethylthiocarbamoyloxy)-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride;
- 40 (±)trans-3,4-dichloro-N-methyl-N-[2-(2,5-dihydro-1H-pyrrol-1-yl)-5-methoxy-1,2,3,4-tetrahydronaphth-1-yl]-benzeneacetamide hydrochloride, and
- 45 (±)trans-3-nitro-N-methyl-N-[2,3-dihydro-2-(pyrrolidin-1-yl)-1H-inden-1-yl]-benzeneacetamide hydrochloride.

22. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering an effective anti-pruritic amount of the composition of claims 19.

5

23. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering an effective anti-pruritic amount of the composition of claims 20.

10

24. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering an effective anti-pruritic amount of the composition of claims 21.

15

25. The method of claim 22 wherein said administration is topical administration.

26. The method of claim 22 wherein said administration is parenteral administration.

27. The method of claim 22 wherein said administration is oral administration.

20

28. The method of claim 22 wherein said administration is rectal administration.

29. A method for the prevention and treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition of claim 20.

25

30. The method of claim 29 wherein said administration is topical administration.

31. The method of claim 29 wherein said administration is parenteral administration.

30

32. The method of claim 29 wherein said administration is oral administration.

33. The method of claim 29 wherein said administration is rectal administration.

35

34. A method for the prevention and treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition of claim 21.


35. The method of claim 34 wherein said administration is topical administration.

36. The method of claim 34 wherein said administration is parenteral administration.

5 37. The method of claim 34 wherein said administration is oral administration.

38. The method of claim 34 wherein said administration is rectal administration.

39. An anti-pruritic pharmaceutical composition comprising a compound of formula
10 III or a pharmaceutically acceptable salt thereof



III

15 wherein
n is 0-1;

R is unsubstituted phenyl or phenyl substituted with one to three substituents selected from the group consisting of halogen, C₁₋₆ alkyl, hydroxy, -O-CO-NH₂, -O-CO-NHalkyl,
20 -O-CO-N(alkyl)₂, C₁₋₆ alkoxy, trifluoromethyl, C₁₋₄-alkoxy-C₁₋₄ alkyloxy, carboxy-C₁₋₄ alkyloxy, nitrile, nitro and amino; or mono or dialkyl amino, amide, sulfonamide, carboxamide; or mono or disubstituted carboxamide, ureido, or mono and di-alkylsubstituted ureido; or

25 R represents an alkyl or cycloalkyl group having up to 7 carbon atoms, wherein the cycloalkyl moiety, where present, can be optionally substituted by one or more substituents selected from the group consisting of from hydroxy, amino, amidino, guanidino, aminocarbonyl, carboxy, C₁₋₆ alkoxy, (C₁₋₆ alkoxy)carbonyl, (C₃₋₆ alkenyloxy)carbonyl, (C₃₋₆ alkynyloxy)carbonyl, C₁₋₆ alkanoyloxy, C₁₋₆ alkylsulfide, C₁₋₆ alkylsulfoxide, C₁₋₆ alkylsulfone, C₁₋₆(monoalkylamino)carbonyl, C₁₋₆ acylamino, C₁₋₆
30 acylmethylamino and C₁₋₆ monoalkylamino; or

R represents the group $-B-R^7$ in which B represents $-CH_2-$, $-CH(CH_3)-$ or a single bond and R^7 represents an optionally substituted C_{6-10} carbocyclic aryl group with one to three substituents selected from the group consisting of halogen, C_{1-6} alkyl, hydroxy, $-O-CO-NH_2$, $-O-CO-NHalkyl$, $-O-CO-N(alkyl)_2$, C_{1-6} alkoxy, trifluoromethyl, C_{1-4} -alkoxy- C_{1-4}

5 alkyloxy, carboxy- C_{1-4} alkyloxy, nitrile, nitro and amino; or mono or dialkyl amino, amide, sulfonamide, carboxamide; mono or disubstituted carboxamide ureido; and mono or di-alkylsubstituted ureido; or

R represents the group $-D-R^8$ in which D represents a single bond, $-CH_2-$, $-CH(CH_3)-$, $-CH_2O-$, $-CH(CH_3)O-$, $-CH_2S-$, $-CH(CH_3)S-$, $-CH_2NH-$ or $-CH(CH_3)NH-$ and R^8 represents
 10 a 4-6 membered heterocyclic ring containing up to 4 heteroatoms selected from the group consisting of oxygen, sulfur and nitrogen, the heterocyclic ring optionally being substituted on nitrogen or sulfur by oxygen or on nitrogen by hydroxy or C_{1-3} alkyl and/or the ring optionally being substituted on carbon by one or more substituents selected from the group consisting of amino, hydroxy, thio (and their tautomers), cyano, halogen, C_{1-3}
 15 alkoxy, C_{1-3} monoalkylamino, C_{1-3} acylamino, C_{1-3} acylmethylamino, and C_{1-3} alkylthio;

R^1 and R^2 are independently selected from the group consisting of H, C_{1-6} alkyl, C_{3-5} alkenyl, C_{3-5} alkynyl, and C_{4-7} cycloalkylalkyl group; or R^2 can be taken together with R^1 and the nitrogen to which they are attached to form a heterocyclic ring which may optionally contain a further heteroatom selected from the group consisting of oxygen,
 20 nitrogen, and sulfur, said heterocyclic ring selected from the group consisting of 1-azetidiny and 1-pyrrolidiny said 1-pyrrolidiny optionally substituted at the 3-position by OH, $-CH_2OH$, tri(C_1-C_6 alkyl)silyloxy, acyloxy, C_{1-6} alkyl, C_{1-6} alkoxy or C_{1-6} alkanoyloxy; 1-piperaziny optionally substituted at the 4-position by alkyl of 1 to 3 carbon atoms; 1-morpholino; 2,5-dihydro-1H-pyrrol-1-yl; 3-azabicyclo[3.1.0]hexan-3-yl;
 25 or 3-azabicyclo[3.2.0]heptan-3-yl;


R^3 represents hydrogen, C_{1-7} alkyl, $-CH_2$ -phenyl or heterocyclic wherein the phenyl or heterocyclic groups may be substituted with one to three substituents selected from the group consisting of halo, C_{1-4} alkyl, C_{1-4} alkoxy and methoxycarbonyl; mono-, di- or tri-halomethyl; cyano; COR^9 , $CH=NOR^{10}$, OR^{10} , SR^{10} , CH_2CN , CH_2OR^{10} , CH_2SR^{10} ,
 30 $CH_2S(O)R^{10}$, $CH_2S(O)_2R^{10}$, $CH_2N(R^{10})R^{11}$, $CH_2(R^{10})R^{11}$, $CH_2NR^{10}OH$, $CH_2N(COR^{10})OH$,

CH₂NR¹⁰COR¹¹, CH₂NR¹⁰S(O)₂R¹¹, or CH₂OCOR¹⁰, wherein R⁹ is hydrogen, hydroxy, amino, NHOH, NHOCH₃, pyridylamino, NHN(CH₃)₂, C₁₋₄ alkoxy, benzyloxy, C₁₋₄ alkylamino, di-C₁₋₄ alkylamino, C₁₋₄ alkyl or C₁₋₄ alkylthio; R¹⁰ and R¹¹ are each hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy or C₇₋₁₁ phenylalkyl), or OR¹², wherein R¹² is hydrogen, C₁₋₄ alkyl or a hydroxy protecting group;

X represents -CO-, or -SO₂-;

Y represents a single bond wherein only one of R⁴-R⁶ is attached, a tetrahedral carbon, -OC-, -SC-, -S(O)C-, -S(O)₂C-, or -CH₂C-;

R⁴, R⁵, and R⁶ are independently selected from the group consisting of hydrogen, hydroxy, alkoxy, C₁₋₄ alkylendioxy, C₁₋₈ cyclic and acyclic alkyl; substituted or unsubstituted carbocyclic aromatic or heterocyclic aromatic group selected from the group consisting of phenyl, naphthyl, biphenyl, indanyl, 1-tetralone-6-yl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl and benzothienyl, each of which may be substituted with one to three substituents selected from the group consisting of halo, cyano, -OCONH₂, -OCONHalkyl, -OCON(alkyl)₂, -OCOalkyl, -NHCHO, -NHCOalkyl, ureido, -NHCONHalkyl, -NalkylCONHalkyl, -NHCON(alkyl)₂, -NalkylCON(alkyl)₂, -NHOSO₂alkyl, -COalkyl, -CONH₂, -CONHalkyl, -CON(alkyl)₂, -CH₂CONH₂, -CH₂CONHalkyl, -CH₂CON(alkyl)₂, -OCH₂CONH₂, -OCH₂CONHalkyl, -OCH₂CON(alkyl)₂, C₁₋₄ alkyl, C₁₋₄ alkoxy, amino, hydroxy, nitro, trifluoromethyl, -SO₂alkyl, -SOalkyl, and mesyl; or R⁵ and R⁶ can together form the following structure



wherein R¹³ and R¹⁴ are independently selected from the group consisting of hydrogen, halogen, hydroxy, alkoxy, mono-, di- or tri-halomethyl, amino, -NHalkyl, -N(alkyl)₂, -NHCOalkyl, ureido, nitro, and methylenedioxy; and

D represents -CH₂-, -O-, -S-, -NH-, -CH₂CH₂-, -CH=CH-, -CH₂NH-, or -CH₂Nalkyl-; in a pharmaceutically carrier.

40. The anti-pruritic pharmaceutical composition of Claim 39 wherein said compound
5 is selected from the group consisting of:

N-methyl-N-{[(1S)-1-phenyl-2-[(3S)-(3-hydroxypyrrolidin-1-yl)]ethyl]-2,2-diphenylacetamide hydrochloride,

3,4-dichloro-N-methyl-N-[(1S)-1-phenyl-2-(1-pyrrolidinyl)ethyl]benzeneacetamide hydrochloride,

- 10 N-methyl-N-{[(1S)-1-phenyl-2-[(3S)-(3-hydroxypyrrolidin-1-yl)]ethyl]-2-aminophenylacetamide hydrochloride,

3,4-dichloro-N-methyl-N-[(1S)-1-isopropyl-2-(1-pyrrolidinyl)ethyl]benzeneacetamide hydrochloride,

- 15 3,4-dichloro-N-methyl-N-[(1S)-1-(O-acetic acid-3-hydroxyphenyl)-2-(1-pyrrolidinyl)ethyl]benzeneacetamide hydrochloride, and

N-methyl-N-[(1S)-1-phenyl-2-(1-pyrrolidinyl)ethyl]-2,2-diphenylacetamide hydrochloride.

41. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-
20 pruritic amount of the composition of claim 39.

42. The method of claim 41 wherein said administration is topical administration.

43. The method of claim 41 wherein said administration is parenteral administration.

- 25 44. The method of claim 41 wherein said administration is oral administration.

45. The method of claim 41 wherein said administration is rectal administration.

46. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition of claim 40.

5 47. The method of claim 46 wherein said administration is topical administration.

48. The method of claim 46 wherein said administration is parenteral administration.


49. The method of claim 46 wherein said administration is oral administration.

10

50. The method of claim 46 wherein said administration is rectal administration.

51. An anti-pruritic pharmaceutical composition comprising a compound of formula IV or a pharmaceutically acceptable salt thereof

15



IV

wherein:

20

R_1 and R_2 are the same or different and are hydrogen, C_{1-6} alkyl, C_{3-6} alkenyl, C_{3-6} cycloalkyl or C_{4-12} cycloalkylalkyl groups, or R_1 and R_2 together form a C_{2-8} branched or linear polymethylene or C_{2-6} alkenylene group, each of which may be optionally substituted with a hetero-atom; or $-NR_1R_2$ forms a 5-membered (optionally containing an oxygen atom adjacent to the nitrogen) or 6-membered ring, which rings optionally contains one unit of unsaturation and which is unsubstituted or substituted with hydroxy, C_{1-6} acyloxy, oxo, methylene, $-COR_{10}$ where R_{10} represents C_{1-6} alkyl, $-OR_{11}$ or $-NHR_{11}$ and R_{11} represents hydrogen, C_{1-6} alkyl, aryl, $Ar(C_{1-6})alkyl$, or $N=NOR_{12}$ (where R_{12} represents C_{1-6} alkyl;

25

R₃ is hydrogen, C₁₋₆ alkyl; or phenyl; or R₃ together with R₁ form a -(CH₂)₃- or -(CH₂)₄- group;

R₄ is C₁₋₆ alkyl, or phenyl;

5

R₅ is hydrogen, or together with R₄ forms a C₂₋₅ linear polymethylene group;

R₆ represents hydroxy, C₁₋₆ alkyl, C₁₋₆ hydroxyalkyl, C₁₋₆ carboxyalkyl, phenyl, oxo, amino, carboxy, amido, -COR₁₃, -CO₂R₁₃ or -COCO₂R₁₃ where R₁₃ represents a hydrogen atom or an unsubstituted or substituted C₁₋₁₀ hydrocarbon moiety; -NR_xCOR_x where R_x represents C₁₋₆ alkyl, optionally substituted methylene or R₆ together with the E atom to which it is attached, forms a 5 or 6-membered ring containing one or more heteroatoms;

10

R₇ is hydrogen, or together with R₆ forms an optionally substituted or unsubstituted single or fused aryl or heterocyclic ring, containing from 5 to 12 ring atoms and comprising up to four heteroatoms in the ring selected from the group consisting of oxygen, nitrogen and sulphur, which may be substituted with hydrogen, C₁₋₆ alkyl, -CH₂OR₁₄, halogen, hydroxy, C₁₋₆ alkoxy, C₁₋₆ alkoxycarbonyl, thiol, C₁₋₆ alkylthio, -OCOR₁₅, -NHCOR₁₆, -NHSO₂R₁₇ or -CH₂SO₂NR₁₈R₁₉, in which each of R₁₄ to R₁₉ is independently hydrogen, C₁₋₆ alkyl, aryl or aralkyl;

20

A is aryl or heteroaryl ring, optionally mono or disubstituted with C₁₋₆ alkyl, C₂₋₆ alkenyl, C₁₋₆ haloalkyl, C₂₋₆ haloalkenyl, C₂₋₆ haloalkynyl, aryl, aralkyl, hydroxy, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, thiol, C₁₋₆ alkylthio, C₁₋₆ haloalkylthio, halogen, nitro, cyano, carboxy, aryloxy, aralkoxycarbonyl, carbamoyl, sulfonyl or sulfamoyl;

25

E represents methylene, sulphur, oxygen or an imino group;

R₈ is hydrogen or C₁₋₆ alkyl; and

30

R₉ is hydrogen or together with R₈ may form the group -(CRaRa)_m-C(=Y)- wherein Ra is hydrogen or C₁₋₆ alkyl having up to a maximum of 3 alkyl groups;

m is 1, 2, or 3; and

- 5 Y represents two hydrogens or oxygen,
in a pharmaceutically acceptable vehicle.

52. The anti-pruritic pharmaceutical composition of claim 51 wherein said compound is selected from the group consisting of:

10

1-(Pyrrolidin-1-yl)methyl-2-(3,4-dichlorophenyl)-acetyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinoline;

15

8-[(3,4-Dichlorophenyl)acetyl]-7-(1-pyrrolidinylmethyl)-1,4-dioxo-8-aza[4.5]spirodecane;

Methyl 4-[3,4-dichlorophenyl)acetyl]-3-(1-pyrrolidinylmethyl)-1-piperazinecarboxylate

20

1-[(3,4-Dichlorophenyl)acetyl]-2-[(3-oxo-1-pyrrolidinyl)methyl]-piperidine.

[S-(RR)]-(-)-5-[(3,4-Dichlorophenyl)acetyl]-4,5,6,7-tetrahydro-4[(3-hydroxy-1-pyrrolidinyl)methyl]furo[3,2-c]pyridine;

25

[S-(RS)]-4-Acetyl-1-[(3,4-dichlorophenyl)acetyl]-2-[(3-hydroxy-1-pyrrolidinyl)methyl]pyridine;

2-[(3,4-Dichlorophenyl)acetyl]-1,2,3,4-tetrahydro-1-(1-pyrrolidinyl)methyl)-5-isoquinolinol;

30

4-(Pyrrolidin-1-yl)methyl-5-(3,4-dichlorophenyl)acetyl-4,5,6,7-tetrahydrothieno[3,2,-c]pyridine;

1-[(5,6-Dichloro-3-oxoindan-1-carbonyl)-2-pyrrolidin-1-ylmethyl]piperidine;

2-(3,4-Dichlorophenyl)acetyl-3-(pyridin-1-yl)methyl-decahydroisoquinoline;

35

1-(4-Trifluoromethylphenyl)acetyl-2-(3-hydroxypyrrolidin-1-yl)methyl-4,4-dimethyl piperidine;

40

4-Acetyl-1-[(3,4-dichlorophenyl)acetyl]-2-[(S)-3-hydroxy-1-pyrrolidinyl)methyl]piperazine;

4-Acetyl-1-[(4-methylsulphonylphenyl)acetyl]-2-[(S)-3-hydroxy-1-pyrrolidinyl)methyl]piperazine;

5 4-(2-Ethanol)-1-[(3,4-dichlorophenyl)acetyl]-2-[(S)-3-hydroxy-1-pyrrolidinyl)methyl]piperazine;

4-Spirohydantoin-1-[(3,4-dichlorophenyl)acetyl]-2-[(pyrrolidinyl)methyl]piperazine; and

10 4-[(S)-3-hydroxy-1-pyrrolidinyl)methyl]-5-[3,4-dichlorophenyl)acetyl]-4,5,6,7-tetrahydroimidazo [4,5-c]pyridine.

53. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition of claim 51.

54. The method of claim 53 wherein said administration is topical administration.

55. The method of claim 53 wherein said administration is parenteral administration.

56. The method of claim 53 wherein said administration is oral administration.

57. The method of claim 53 wherein said administration is rectal administration.

58. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition of claim 52.


59. The method of claim 58 wherein said administration is topical administration.

60. The method of claim 58 wherein said administration is parenteral administration.

61. The method of claim 58 wherein said administration is oral administration.

62. The method of claim 58 wherein said administration is rectal administration.

63. An anti-pruritic pharmaceutical composition comprising a compound of formula V or a pharmaceutically acceptable salt thereof



V

wherein

----- represents a single or double bond;

R^1 represents an alkyl group having 1-5 carbon atoms, a cycloalkylalkyl group having 4-7 carbon atoms, a cycloalkenylalkyl group having 5-7 carbon atoms, an aryl group having 6-12 carbon atoms, an aralkyl group having 7-13 carbon atoms, an alkenyl group having 4-7 carbon atoms, an allyl group, a furan-2-ylalkyl group having 1-5 carbon atoms, or a thiophen-2-ylalkyl group having 1-5 carbon atoms;

R^2 represents a hydrogen atom, a hydroxy group, a nitro group, an alkanoyloxy group having 1-5 carbon atoms, an alkoxy group having 1-5 carbon atoms, an alkyl group having 1-5 carbon atoms, or $-NR^9R^{10}$ wherein R^9 represents a hydrogen atom or an alkyl group having 1-5 carbon atoms, and R^{10} represents a hydrogen atom; an alkyl group having 1-5 carbon atoms, or $-C(=O)R^{11}$ wherein R^{11} represents a hydrogen atom, a phenyl group or an alkyl group having 1-5 carbon atoms;




R^3 represents a hydrogen atom, a hydroxy group, an alkanoyloxy group having 1-5 carbon atoms, or an alkoxy group having 1-5 carbon atoms;







A represents $-XC(=Y)-$, $-XC(=Y)Z-$, $-X-$, $-XSO_2-$, or $-OC(OR^4)R^4-$ where, X, Y and Z each independently represent NR^4 , S or O wherein R^4 represents a hydrogen atom, a straight-chain or branched chain alkyl group having 1-5 carbon atoms or an aryl group having 6-12 carbon atoms, and wherein R^4 may be identical or different;

B represents a valence bond, a straight-chain or branched chain alkylene group having 1-14 carbon atoms which may be substituted with at least one substituent selected from the group consisting of an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, a hydroxy group, fluorine, chlorine, bromine, iodine, an amino group, a nitro group, a cyano group, a trifluoromethyl group and a phenoxy group, and wherein 1 to 3 methylene groups may be replaced with carbonyl groups, an acyclic unsaturated hydrocarbon containing from 1 to 3 double bonds and/or triple bonds and having 2-14 carbon atoms which may be substituted with at least one substituent group selected from the group consisting of an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, a hydroxy group, fluorine, chlorine, bromine, iodine, an amino group, a nitro group, a cyano group, a trifluoromethyl group and a phenoxy group, and wherein from 1 to 3 methylene groups may be replaced with carbonyl groups, or a straight-chain or branched chain saturated or unsaturated hydrocarbon group containing from 1 to 5 thioether, ether and/or amino bonds and having 1-14 carbon atoms wherein hetero atoms are not bonded directly to A, and 1 to 3 methylene groups may be replaced with carbonyl groups;

R⁵ represents a hydrogen atom or an organic group (which may be substituted with at least one or more substituent groups selected from the group consisting of an alkyl group having 1-5 carbon atoms, an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, a hydroxy group, fluorine, chlorine, bromine, iodine, an amino group, a nitro group, a cyano group, an isothiocyanate group, a trifluoromethyl group and a methylenedioxy group); or

R₅ is





wherein

Q is N, O or S;

T is CH, N, S or O;

5 l is 0-5;

m and n are ≥ 0

m + n ≤ 5 ;

R⁶ represents a hydrogen atom;

10

R⁷ represents a hydrogen atom, a hydroxy group, an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, or R⁶ and R⁷ together represent -O-, -CH₂- or -S-;


15 R⁸ represents a hydrogen atom, an alkyl group having 1-5 carbon atoms, or an alkanoyl group having 1-5 carbon atoms in a pharmaceutically acceptable carrier.

64. The anti-pruritic pharmaceutical composition of claim 63 wherein

20 R¹ is an alkyl group having 1-5 carbon atoms, a cycloalkylmethyl group having 4-7 carbon atoms, a cycloalkenylmethyl group having 5-7 carbon atoms, a phenylalkyl group having 7-13 carbon atoms, an alkenyl group having 4-7 carbon atoms, an allyl group, a furan-2-yl-alkyl group having 1-5 carbon atoms and a thiophen-2-yl-alkyl group having 1-5 carbon atoms;

R^2 is hydrogen, hydroxy, nitro, acetoxy, methoxy, methyl, ethyl, propyl, amino, dimethylamino, acetylamino or benzoylamino groups; or

5 R^4 is



Formula V-1

wherein

Q is N, O or S;

10 T is CH, N, S or O;

m and n are ≥ 0 and

$m + n \leq 5$;

B is $-(CH_2)_n-$ wherein $n = 0-6$, $-(CH_2)_n-C(=O)-$ wherein $n = 1-4$, $-CH=CH-(CH_2)_n-$

wherein $n = 0-4$, $-C\equiv C-(CH_2)_n-$ wherein $n=0-4$, $-CH_2-O-$, $-CH_2-S-$, $-CH_2-O-(CH_2)_2-O-$

15 $(CH_2)_2-$, $-CH_2-O-CH_2-NH-CH_2-O-CH_2-$ and $-CH_2-O-CH_2-S-CH_2-O-CH_2-$;

R^5 is hydrogen or an organic group of Formula V-1 said organic group may be substituted with at least one substituent group selected from the group consisting of an alkyl group having 1-5 carbon atoms, an alkoxy group having 1-5 carbon atoms, an alkanoyloxy group having 1-5 carbon atoms, a hydroxy group, fluorine, chlorine, bromine, an amino group, a nitro group, a cyano group, an isothiocyanate group and a trifluoromethyl group, in a pharmaceutically acceptable carrier.

20

65. The anti-pruritic pharmaceutical composition of claim 64 wherein

R^1 is methyl, ethyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclopentenylmethyl, cyclohexenylmethyl, benzyl, phenethyl, trans-2-butenyl, 2-methyl-2-butenyl, allyl, furan-2-yl-methyl or thiophen-2-yl-methyl;

R^2 is hydrogen, hydroxy, nitro, acetoxy, methyl or dimethylamino;

- 5 R^3 is $-NR^4C(=O)-$, $-NR^4C(=S)-$, $-NR^4C(=O)O-$, $-NR^4C(=O)NR^4-$, $-NR^4C(=S)NR^4-$ or $-NR^4SO_2-$;

R^4 is a straight-chain or branched alkyl group having 1-5 carbon atoms;

B is $-(CH_2)_n-$ wherein $n=0-6$, $-CH=CH(CH_2)_n-$ wherein $n=0-4$, $-C\equiv C-(CH_2)_n-$ wherein $n=0-4$, $-CH_2-O-$ or $-CH_2-S-$; and

- 10 R^5 is hydrogen, phenyl, 3,4-dichlorophenyl, 4-chlorophenyl, 3-chlorophenyl, 3,4-difluorophenyl, 4-fluorophenyl, 3-fluorophenyl, 2-fluorophenyl, 4-bromophenyl, 3-bromophenyl, 2-bromophenyl, 4-nitrophenyl, 3-nitrophenyl, 2-nitrophenyl, 4-trifluoromethylphenyl, 3-trifluoromethylphenyl, 2-trifluoromethylphenyl, 4-methylphenyl, 3-methylphenyl, 2-methylphenyl, 4-methoxyphenyl, 3-methoxyphenyl, 2-methoxy, 3-furanyl, 2-furanyl, 3-thienyl, 2-thienyl, cyclopentyl or cyclohexyl, in a
15 pharmaceutically acceptable carrier.

66. The anti-pruritic pharmaceutical composition of claim 65 wherein said compound is selected from the group consisting of:

- 20 17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-3-phenylpropionamido)morphinan;
- 25 17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-trans-3-(3-furyl)acrylamido)morphinan;
- 17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-trans-3-cyclohexylacrylamido)morphinan;
- 30 17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-trans-3-(4-trifluoromethylphenyl)acrylamido)morphinan;

17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 α -(N-methyl-trans-3-(3-thiophenyl)acrylamido)morphinan;

- 5 17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-trans-3-phenylacrylamido)morphinan;

17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-trans-2-hexenamido)morphinan; and

10

17-cyclopropylmethyl-4,5 α -epoxy-3,14 β -dihydroxy-6 β -(N-methyl-phenylpropiolamido)morphinan

67. A method for the prevention or treatment of pruritus in a mammal in need of such
15 prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition of claim 63.

68. The method of claim 67 wherein said administration is topical administration.

20

69. The method of claim 67 wherein said administration is parenteral administration.

70. The method of claim 67 wherein said administration is oral administration.

71. The method of claim 67 wherein said administration is rectal administration.

25

72. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition of claim 64.

30

73. The method of claim 72 wherein said administration is topical administration.

74. The method of claim 72 wherein said administration is parenteral administration.

75. The method of claim 72 wherein said administration is oral administration.

35

76. The method of claim 72 wherein said administration is rectal administration.

77. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition of claim 65.

40

78. The method of claim 77 wherein said administration is topical administration.

79. The method of claim 77 wherein said administration is parenteral administration.

80. The method of claim 77 wherein said administration is oral administration.

81. The method of claim 77 wherein said administration is rectal administration.

82. A method for the prevention or treatment of pruritus in a mammal in need of such prevention or treatment comprising administering to said mammal an effective anti-pruritic amount of the composition of claim 66.

83. The method of 82 wherein said administration is topical administration.

84. The method of claim 82 wherein said administration is parenteral administration.

85. The method of claim 82 wherein said administration is oral administration.

86. The method of claim 82 wherein said administration is rectal administration.

ABSTRACT

Anti-pruritic pharmaceutical formulations containing kappa agonist compounds and methods of prevention or treatment of pruritus in a mammal with the anti-pruritic
5 formulations.

DECLARATION AND POWER OF ATTORNEY FOR UNITED STATES PATENT APPLICATION

X Original Supplemental Substitute

As a below-named inventor, I hereby declare that:

My residence, citizenship and post office address are given below under my name.

I believe I am an original, first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled:

KAPPA AGONIST ANTI-PRURITIC PHARMACEUTICAL FORMULATIONS AND METHOD OF TREATING PRURITUS THEREWITH

the specification of which

X is attached hereto.

_____ was filed on _____ as
Application Serial No. _____
and with amendments through _____ (if applicable).

I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge my duty to disclose information of which I am aware which is material to the examination of this application in accordance with Section 1.56(a) of Title 37 of the Code of Federal Regulations.

I hereby claim foreign priority benefit under Section 119 of Title 35 of the United States Code of any foreign application(s) for patent or inventor's certificate or of any PCT application(s) designating at least one country other than the United States identified below and also identify below any foreign application for patent or inventor's certificate or any PCT application(s) designating at least one country other than the United States filed by me on the same subject matter and having a filing date before that of the application(s) from which priority is claimed:

Country	Number	Filing Date	Priority Claimed	
			Yes	No

I hereby claim benefit under Section 120 of Title 35 of the United States Code of any United States application(s) or PCT application(s) designating the United States identified below and, insofar as the subject matter of each of the claims of this application is not disclosed in said prior application(s) in the manner provided by the first paragraph of Section 112 of Title 35 of the United States Code, I

acknowledge my duty to disclose material information of which I am aware as defined in Section 1.56(a) of Title 37 of the Code of Federal Regulations which occurred between the filing date of the prior application(s) and the national or PCT filing date of this application:

Application Serial No. Filing Date Status

I hereby appoint Imre Balogh, Reg. No. 26,835 as my attorney with full power of substitution and revocation to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.

SEND CORRESPONDENCE TO:

276 Smith School Road
Perkasie, PA 18944

DIRECT TELEPHONE CALLS TO:

Imre Balogh
IMRE (JIM) BALOGH

Telephone No. (610) 889-8995

I hereby declare that all statements made herein and in the above-identified specification of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of first joint inventor John J. Farrar
Inventor's signature [Signature] Date 7/10/97
Residence 1066 Bodine Road, Chester Springs, Pennsylvania 19425
Post Office Address 1066 Bodine Road, Chester Springs, Chester County, Pennsylvania 19425
Citizenship United States

Full name of second joint inventor An-Chih Chang
Inventor's signature [Signature] Date 7/10/97
Residence 1447 Neshaminy Valley, Bensalem, Pennsylvania 19020
Post Office Address 1447 Neshaminy Valley, Bensalem, Pennsylvania 19020
Citizenship Costa Rica

Full name of third joint inventor Virendra Kumar
 Inventor's signature Virendra Kumar Date 7/10/97
 Residence 14 Lynne Circle, Paoli, Chester County, Pennsylvania 19301
 Post Office Address 14 Lynne Circle, Paoli, Chester County, Pennsylvania 19301
 Citizenship United States

Full name of fourth joint inventor Wei Yuan Zhang
 Inventor's signature Wei Yuan Zhang Date 7-10-97
 Residence 154 Hunsicker Drive, Collegeville, Montgomery County, Pennsylvania 19426
 Post Office Address 154 Hunsicker Drive, Collegeville, Montgomery County, Pennsylvania 19426
 Citizenship United States

Full name of eighth joint inventor Alan Cowan
 Inventor's signature Alan Cowan Date 6/27/97
 Residence 549 Melissa Drive, Ambler, Montgomery County, Pennsylvania 19002
 Post Office Address 549 Melissa Drive, Ambler, Montgomery County, Pennsylvania 19002
 Citizenship British